WALNUT CREEK SUPPLY

THE COMPLETE CARE GUIDE

Veiled Chameleon

Hydration, gut-loading and the lighting that keeps a chameleon alive

Family-owned in Christiana, Lancaster County, Pennsylvania

10% of every purchase supports no-kill animal shelters nationwide

14 sections · walnutcreeksupply.com

CONTENTS

What's Inside

01	Quick Start: Your First 48 Hours
02	Meet Your Veiled Chameleon
03	Enclosure Setup & Layout
04	Substrate Guide
05	Heat, UVB & Humidity — The 3 Numbers
06	Feeding & Gut-Loading
07	Hydration — The Critical Section
08	Shedding
09	Common Challenges & Solutions
10	Handling
11	Routine Care Calendar
12	When to See a Vet
13	FAQ
14	Every Purchase Gives Back

New owner? *Go straight to Section 1. It is written so you can set up a safe, correct home and get through the first 48 hours without reading anything else first.*

Veiled chameleons are among the most visually spectacular reptiles a keeper can own — and among the most demanding. That combination is exactly why so many end up in rescue collections within their first year. This guide does not exist to talk you out of a veiled chameleon. It exists to tell you the truth about what one actually needs, delivered by a family in Christiana, Lancaster County, Pennsylvania who believe that any animal entering a home deserves a keeper who went in with eyes wide open. If you have researched the ventilation requirements, set up the misting system, located an exotic vet, and are genuinely prepared for a reptile that communicates almost exclusively in subtle physical signals — then you are exactly the person this guide was written for. The reward is real. A healthy, settled veiled chameleon on a well-planted screen enclosure, firing up bright greens and yellows as he basks in the morning sun, is one of the most extraordinary animals you will ever share a home with. But the baseline has to be right from day one.

SECTION 1

Quick Start: Your First 48 Hours

1. **Build the entire enclosure before any animal travels toward your home.** A minimum **24" × 24" × 48" tall screen enclosure** — keyword: screen. Glass terrariums are not acceptable housing for chameleons under any circumstances. The enclosure should be vertical, not horizontal. Have it fully assembled, planted with branches and live non-toxic plants, and running with all equipment calibrated before pickup or delivery day.
2. **Install and test the misting system first.** A chameleon arriving into a dry enclosure on its first morning is a chameleon already in deficit. Program the mister to run at least morning and evening, long enough to coat every leaf surface with visible droplets. Confirm water drains away completely between cycles — chameleons need wet-dry cycling, not standing dampness.
3. **Establish the temperature gradient and confirm it with real measurements.** Basking branch surface **85–95°F** (measure the exact surface under the basking lamp with an infrared temperature gun), ambient warm side air **75–80°F**, cool end **68–72°F**. An unverified basking spot is not a basking spot — it is a guess.
4. **Install UVB over the basking zone.** A T5 HO 5.0 or 6% tube (not 10% — that is too intense for an arboreal forest species) positioned so the chameleon basks within 6 to 12 inches of the lamp. Photoperiod: 12 hours on, 12 hours off, on a timer.
5. **Plant the enclosure densely.** Live pothos, hibiscus, or schefflera not only provide visual security — they hold water droplets the chameleon will drink, and veiled chameleons will also graze plant leaves for supplemental hydration and trace nutrition. All plants must be confirmed non-toxic, soil must be covered or potted to prevent ingestion, and no fertilizer or pesticide residue should be present. Rinse ficus benjamina leaves thoroughly before introducing; the sap can irritate a chameleon's mouth.
6. **Source feeder insects and gut-load them before the chameleon arrives.** Dubia roaches or crickets that have been fed leafy greens and a quality dry gut-load blend for 24 to 48 hours are what your chameleon needs on arrival. Insects with empty guts provide almost no nutrition. Prepare this in advance, not the morning you pick up the animal.
7. **Find an exotic animal vet who sees chameleons specifically before the animal enters your home.** Chameleons mask illness until they are critically ill. A newly arrived chameleon who shows no symptoms on day one can be in serious trouble by day four. Calling around after a problem emerges wastes time you do not have. Tape the number to the enclosure.
8. **Place the chameleon in the enclosure, offer a light misting within the first two hours, and then step away.** Do not handle a newly arrived chameleon. Do not stare at it. Do not tap the enclosure. Move slowly and quietly around the room. A chameleon who is gaping (holding the mouth open while at rest), rocking side-to-side, or turning very dark immediately after introduction is telling you something is wrong — revisit temperatures, confirm no drafts or stress sources, and call the vet if symptoms persist through the first day.

9. **Do not attempt handling until the chameleon has eaten voluntarily at least three to five times.** Chameleons tolerate handling rather than enjoying it. The relationship between a keeper and a settled chameleon is built over weeks, not days.
10. **Set a calendar reminder for six months out to replace the UVB bulb.** Unlike most reptile species, chameleons are in Ferguson Zone 3 — they receive high UV exposure in the wild and need a lamp at full output to match it. UVB output drops significantly before the lamp visibly dims. Every 6 months, not annually. Write it down now.

First-Time Starter Bundle

Everything that should be purchased, assembled, and tested before any chameleon comes home.

Essentials

- **A tall screen enclosure, minimum 24" × 24" × 48"** — all-screen or mesh-paneled construction; no solid glass walls; designed for full cross-ventilation on multiple sides. PVC-framed screen enclosures purpose-built for chameleons are widely available and purpose-designed for drainage. This is the single most important hardware choice in the entire setup.
- **A basking lamp** — an incandescent or halogen bulb positioned above the top screen to create a hotspot on a horizontal branch at **85–95°F** at chameleon level. The wattage depends on ambient room temperature; start at 50W, measure with an infrared gun, and adjust rather than assuming.
- **A T5 HO UVB fluorescent tube, 5.0 or 6%** — positioned over the basking zone inside the enclosure hood or on top of the screen. Replace every 6 months without exception. The 10.0 version is too intense for a forest arboreal species spending extended hours near the lamp and has been linked to photokeratoconjunctivitis (UV-induced eye inflammation) in chameleons.
- **Two digital thermometers with probes** — one at basking branch level, one at the cool end of the enclosure. An infrared temperature gun is also essential for direct surface readings.
- **A programmable automatic misting system** — the Mistking Starter, Aquazamp, or equivalent. A hand mister used once a day is not sufficient; automated misting ensures the chameleon receives water when you are at work, asleep, or otherwise occupied. Place a drain tray underneath the enclosure.
- **A drip system or supplemental dripper** — even with an automated mister, a slow drip running for one to two hours each day encourages drinking behavior. A clean repurposed IV bag or a perforated cup positioned to drip onto a branch is a functional low-cost option alongside a commercial dripper.
- **Live, non-toxic plants** — pothos (*Epipremnum aureum*), umbrella plant (*Schefflera arboricola*), hibiscus (*Hibiscus rosa-sinensis*), or ficus benjamina (washed). Quantity matters: the enclosure should look like a small jungle, not a display case with one vine. Dense coverage provides security, multiple basking angles, and extensive leaf surfaces for drinking.
- **Horizontal and diagonal hardwood branches** — natural (untreated) or high-quality artificial, multiple diameters, placed at various heights. One thick branch at the primary basking zone, 6 to 12 inches below the basking lamp. Additional branches at mid-enclosure and lower levels creating a gradient of temperature and light with connected paths between them.
- **A digital hygrometer** — positioned at mid-enclosure. Target 50–70%; expect natural cycling with highs immediately after misting and lows before the next cycle.
- **A feeder insect supply** — dubia roaches and crickets as staples; hornworms and silkworms as occasional variety feeders. Source from a reputable live feeder supplier, never from the wild.
- **Calcium supplement (without D3)** — for dusting feeders at every meal.
- **Calcium with D3** — for twice-monthly supplemental dusting.
- **Reptile multivitamin** — for once-monthly supplemental dusting.

> ***Ready to set up?*** *Browse the essentials in our collections: Reptile and Amphibian Enclosures and Reptile and Amphibian Terrarium Supplies & Accessories.*

Enrichment

- **Pothos vine lengths woven through the branches** — pothos is the hardiest, most adaptable chameleon enclosure plant and nearly impossible to kill under misting conditions. Route lengths along the back and sides for dense visual coverage and additional drinking surfaces throughout the enclosure.
- **Hanging vines, artificial or real** — creates pathway variety between height levels and fills the visual space so the chameleon always has cover within reach. Veiled chameleons feel exposed when there is open vertical space between them and the nearest cover.
- **A secure, smooth-sided feeder cup** attached to a branch — a small cup zip-tied to a mid-enclosure branch keeps crickets or black soldier fly larvae visible and accessible without allowing them to escape into hard-to-reach areas. Chameleons are visual hunters and will reliably target a known feeder station once they learn it.
- **Cork bark rounds or flats** mounted vertically or diagonally as additional resting surfaces and visual barriers.
- **A solid visual barrier on at least one side of the enclosure** — a dark background panel on the rear prevents the chameleon from seeing its own reflection, spotting other animals through the opposite wall, or being stressed by activity in the room. Position the enclosure against a wall if possible and away from high-traffic pathways.
- **A sturdy travel carrier** lined with a diagonal branch and a small rooted pothos cutting for vet visits — a chameleon traveling in a bare plastic box with nothing to grip is a stressed chameleon before it gets to the clinic.

> ***Enrichment, one click away.*** *Shop Reptile and Amphibian Accessories.*

SECTION 2

Meet Your Veiled Chameleon

Wild veiled chameleons (*Chamaeleo calyptratus*) originate from the highland escarpments and forested wadis of Yemen and southwestern Saudi Arabia — specifically the terraced mountain slopes of the Asir and Sarawat ranges, at elevations that can reach nearly 10,000 feet. These are not desert animals. Their homeland is humid, seasonally variable highland scrub and cultivated agricultural terracing, where temperatures drop sharply at night, morning mists roll in off the Red Sea coast, and midday sun can be intense at altitude. This origin story explains almost everything about what captive veiled chameleons require: vigorous ventilation, dramatic day-to-night temperature swings, humidity that cycles rather than stays constantly high, and elevated UVB exposure from genuine highland sunlight.

Trait	Detail
Common name	Veiled chameleon, Yemen chameleon
Scientific name	*Chamaeleo calyptratus*
Native range	Yemen, southwestern Saudi Arabia; highland terraces and forested escarpments
Wild habitat	Mountain scrub, terraced farmland, forested wadis; up to ~3,000 m elevation
Adult length — male	18–24 inches total (snout to tail tip)
Adult length — female	10–14 inches total
Typical lifespan — male	**5–8 years** with good care
Typical lifespan — female	**3–5 years** (egg production is physiologically costly)
Activity pattern	Diurnal — active during daylight hours; sleeps motionless on branches at night
Social structure	**Strictly solitary** — visual contact with another chameleon causes chronic stress
Temperament	Defensive, stress-sensitive; tolerates calm handling; does not seek contact
Relative difficulty	The most beginner-accessible chameleon species — but "beginner chameleon" is not "beginner reptile"

Anatomy you should know

Casque: the helmet-like bony crest on top of a veiled chameleon's head. In adult males, it is tall, prominently ridged, and laterally compressed — the feature that gives the species its common name, as it resembles a veil raised above the face. Females develop a shorter, less prominent casque. The casque channels rainwater and morning dew down the face toward the mouth, a drinking behavior deeply ingrained in the species and the reason why chameleons drink from surfaces rather than from standing pools.

Turret eyes: chameleons have laterally compressed eyeballs set inside cone-shaped, independently mobile turrets — each eye moves and focuses entirely independently of the other, giving the animal close to 360-degree vision without turning its head. When both eyes converge forward on a single target, the chameleon is in the final stages of locking onto prey or a threat. Watch for binocular convergence during feeding sessions — it precedes a tongue strike by a fraction of a second.

Prehensile tail: the tail functions as a fifth limb, coiling around branches to stabilize the body during movement, extension of the tongue, and resting. Unlike a leopard gecko's tail, a chameleon's tail does not drop as a defensive reflex (it is not autotomic) and is not a fat reserve. It is a structural tool, and an injured or kinked tail can compromise the animal's ability to navigate its enclosure.

Zygodactyl feet: each foot has toes grouped into two opposing bundles — two toes opposed against three — that clamp around branches with a grip that is almost mechanical in its precision. This is a complete specialization for arboreal life, and it means chameleons are genuinely poor walkers on flat surfaces. Handle a chameleon with branches or perches available; placing one on an open flat surface causes distress.

Projectile tongue: typically one and a half to two times body length, launched from a hyoid bone accelerator mechanism at extraordinary speed — strikes measured in some chameleon species at under 0.1 seconds. The tongue tip carries a thick adhesive mucus that traps prey on contact. A chameleon that has stopped hunting — tracking feeders with its eyes but not firing the tongue — is a chameleon telling you something is wrong.

Color communication

The popular understanding that chameleons change color for camouflage is partially accurate but incomplete as a working model for a keeper. Color in veiled chameleons is primarily a thermoregulation and communication system.

Healthy basking colors: bright greens with pale yellow or cream banding and occasionally turquoise lateral highlights. A chameleon showing this palette is warming up successfully and feels secure in its environment.

Stress or threat display: dark brown to near-black base coloration with high-contrast yellow, orange, or white spots. Briefly during handling or when the chameleon spots something alarming, this is a normal defensive display. Sustained dark coloration outside of these short windows — particularly at rest in a warm, correctly heated enclosure — is a flag that something is chronically wrong.

Cold or early illness: a uniformly dull, dark brown coloration without the high-contrast spots of the threat display. A dark animal trying to absorb heat will typically orient itself broadside to the basking lamp; if the temperature is correct and the animal is still uniformly dark, investigate health and environment.

Male display coloration: intensely vivid greens, yellows, blues, and turquoise — a palette that can startle new keepers because the contrast is so striking. This is a good sign. It typically appears when a male is at peak body temperature, in good health, and feeling territorial or alert.

The Panther Chameleon as an alternative

The **Panther Chameleon** (*Furcifer pardalis*) is worth mentioning for any keeper researching their first chameleon. Male Panthers are among the most colorful reptiles on earth, with locale-specific morphs that range from blue and red to green and orange. Panthers tend to be slightly more even-tempered than Veileds and have parallel environmental requirements — screen enclosure, misting, UVB, insect diet, night temperature drop — but generally prefer slightly higher humidity (60–80%) and do not graze live plants in the enclosure the way Veileds characteristically do. If a less plant-dependent setup is appealing, Panthers are a reasonable alternative. Neither species is a low-effort animal.

What surprises new owners

The first surprise is the drinking behavior. A new keeper who sets up a beautiful enclosure with a water bowl and no misting system will come home to a dehydrated chameleon. Chameleons in captivity will frequently ignore standing water entirely because their instinct and anatomy are designed to drink from droplets in motion or from wet leaf surfaces after a rain event. The misting system is not optional equipment — it is the water supply.

The second surprise is how fast a sick chameleon deteriorates once symptoms become visible. A chameleon who seems completely normal on Monday morning may be critically ill by Thursday. This is not because chameleons are uniquely fragile on a cellular level — it is because they are hardwired to conceal weakness from predators. By the time sunken eyes, gaping, or rapid color loss are visible, the underlying problem has been developing for a while. Keepers who check in briefly once a day from across the room miss the early window entirely. Daily close observation — actually looking at the animal's eyes, posture, and coloration — is part of the job.

The third surprise is the lifespan, which is short compared to most reptiles a keeper might have owned before. A healthy male veiled chameleon lives 5 to 8 years. A healthy female lives 3 to 5 years — shorter, because egg production, which happens even in females who have never been near a male (they produce infertile clutches routinely), is metabolically demanding and depletes calcium and body reserves with each clutch. Overbreeding and calcium depletion are among the most common causes of premature female death in captive collections.

SECTION 3

Enclosure Setup & Layout

Screen only — not a preference, a requirement

A glass terrarium cannot house a veiled chameleon safely. This is not a matter of preference or budget — it is a biological reality. Glass traps heat, stagnant air, and moisture against the walls. In a glass enclosure with a chameleon inside and a misting cycle running, the ventilation necessary to prevent respiratory infections simply cannot happen. The warm, stagnant, constantly humid microenvironment that results is an ideal culture medium for the bacteria that cause respiratory disease. Chameleons have small tidal volumes and breathe rapidly — they are highly sensitive to poor air quality. The situation compounds: glass also prevents a proper wet-dry humidity cycle from occurring, and stagnant high humidity without air movement is exactly what precipitates infection.

The enclosure must be primarily screen or mesh on multiple faces. PVC-framed all-screen enclosures purpose-built for chameleons are the standard recommendation. The minimum size for one adult is **24" × 24" × 48" tall**.

Why height matters more than floor area

Chameleons are arboreal — they live at height and experience the ground as exposed, threatening space. A properly tall enclosure allows a natural vertical temperature gradient: the basking zone at the top near the lamp, cooler and shadier zones progressively lower, with the floor being the coolest and least-used level. A chameleon in a correctly arranged tall enclosure thermoregulates by moving up and down, exactly as it would in a tree. A low, wide enclosure forces the animal into a single temperature band and provides no meaningful escape to a cooler zone without descending to an area it instinctively avoids.

Branch and plant layout — described so it can be drawn

Beginning at the top of a 48-inch enclosure and working downward:

Top zone (top 12 inches): one thick horizontal basking branch running parallel to the UVB lamp and basking light source — this is the primary morning perch and the location the chameleon will return to each day at first light. The basking surface at this branch should read **85–95°F**. A second slightly thinner branch a few inches below and laterally offset provides an alternative position in the same warm zone.

Upper mid zone (12–24 inches from top): additional diagonal branches crossing the enclosure at angles, anchor points for hanging pothos lengths, and one dense plant cluster near the back providing a visual retreat and a shaded rest area. Ambient temperature here should be **75–80°F**.

Lower mid zone (24–36 inches from top): a schefflera canopy, a routed section of pothos, and branch intersections providing dense coverage. Ambient temperature here **70–75°F** — a meaningful cool retreat for midday shelter.

Bottom zone (bottom 12 inches): drainage layer, potted plant bases, misting nozzle outlet points, and the drip collection tray. Chameleons rarely descend to the floor voluntarily except when a female is preparing to lay eggs. This zone is primarily about waste management, plant root health, and drainage.

Misting nozzle placement: direct at least one nozzle toward upper and mid-level leaf clusters — not only at walls. The goal is leaf surfaces loaded with visible water droplets that persist for 10 to 20 minutes after the misting cycle ends.

Visual barriers: a solid dark background panel on the rear wall. Do not place the enclosure near other reptile enclosures, windows with outdoor animal activity (birds especially), mirrors, or any reflective surface. A chameleon with 340-degree vision that spots a rival — or its own reflection — through the enclosure wall experiences a territorial threat it cannot resolve, and that unresolvable chronic threat is one of the most damaging long-term stressors in captive chameleons.

SECTION 4

Substrate Guide

The floor of a tall screen chameleon enclosure requires different thinking than a terrestrial reptile setup. Because the enclosure receives multiple misting cycles daily, the priority is drainage, plant root health, and waste management rather than thermal mass or burrowing capacity.

Floor option	Best for	Performance	Notes
Drainage layer: LECA (horticultural clay pebbles), 2–3 inches deep	All screen enclosure setups	Excellent drainage; separates pooled water from organic matter above; prevents anaerobic rot beneath the substrate	Separate from the substrate layer above with a fine mesh screen divider to prevent mixing over time
Top layer: organic potting soil + coarse sand, 70:30	Naturalistic setups with deeply rooted live plants	Supports live plant root systems; holds light moisture without waterlogging above the drain layer	Use additive-free potting soil only; no large perlite pellets that could be ingested; depth 2–3 inches above drainage layer
Top layer: coconut coir, rehydrated	Simpler setups where easy replacement matters	Clean, low-dust, holds light moisture; moderate plant support for potted plants sitting in the layer	Replace every 6–8 weeks before the layer becomes anaerobic under regular misting; not ideal for large root systems
Paper towel over drainage layer	Hospital, quarantine, sick animals	Easiest waste inspection; zero particulate ingestion risk; complete hygiene control	Not a permanent setup option; use for newly arrived animals during the first 2–4 weeks or during illness recovery
Potted plants on elevated stands, bare drainage layer floor	Clinical setups prioritizing maximum cleanliness	Maximum drainage; pots are individually removable for cleaning; easiest floor to disinfect	Less natural in appearance; fully functional; a practical choice for keepers who want simple maintenance
Sand alone	**Avoid**	Poor drainage under misting; anaerobic rot risk; ingestion hazard	Inappropriate for any chameleon enclosure at any life stage
Reptile carpet	**Avoid**	Traps waste between fibers; harbors bacteria under a twice-daily misting schedule; cannot drain properly	Inappropriate for misted arboreal setups

For live plant integration: pot each plant individually in its original nursery container with a drainage hole, place pots in the bottom zone, and route branches and vines up through the enclosure so only the plant body is visible. Cover bare soil in each pot with a layer of smooth river pebbles large enough that the chameleon cannot swallow them — this discourages soil digging and inadvertent ingestion.

SECTION 5

Heat, UVB & Humidity — The 3 Numbers

Veiled chameleons require all three environmental parameters to be correct simultaneously and to interact correctly with each other. Specifically, the humidity must cycle upward after misting and then fall back as the enclosure air exchanges — and this requires the ventilation that only a screen enclosure can deliver. A correct humidity cycling pattern in a glass enclosure is physically impossible. Every chronic health problem in a captive veiled chameleon traces back to at least one of these three parameters being persistently off.

Parameter	Target value	How to measure it	What goes wrong when it's off
Basking spot surface temperature	85–95°F	Infrared temperature gun pointed at the basking branch surface	Too high: thermal burns, active avoidance of the basking zone, failure to thermoregulate; too low: poor digestion, immune suppression, chronic inactivity
Ambient warm-side air temperature	75–80°F	Probe thermometer at upper mid zone height	Below 72°F: incomplete thermoregulation; above 85°F ambient: heat stress, immune function decline
Cool end air temperature	68–72°F	Probe thermometer at lower enclosure level	Too warm: no functional gradient exists; too cool: hypothermia risk below 60°F
Nighttime temperature drop	55–65°F (target after lights out)	Same probes with lights off overnight	No night drop: disrupts circadian rhythm, suppresses long-term appetite and health; below 50°F: hypothermia risk
Relative humidity — post-misting peak	70–80%	Digital hygrometer at mid-enclosure during or immediately after misting	Too low at peak: inadequate drinking opportunity on leaf surfaces; inadequate shed moisture
Relative humidity — midday trough	30–50%	Same hygrometer mid-cycle, between misting sessions	Staying above 60–70% all day without air movement through screen: respiratory infection risk
UVB — Ferguson Zone 3	UVI 1.0–2.6	Solarmeter 6.5R; or T5 HO 5.0/6% tube at 6–12 inches from basking branch	Too low: metabolic bone disease (MBD) risk, reduced appetite, poor immune function; too high (Zone 4+): photokeratoconjunctivitis, eye damage

The night-drop protocol

The highland origin of veiled chameleons means they evolved under significant day-to-night temperature swings. A room that stays at 72–75°F all night — comfortable for a human — is too warm for a veiled chameleon's long-term health. The target **55–65°F overnight range** is cooler than most reptile guides recommend and cooler than most new keepers expect.

In practice, many homes cool sufficiently in autumn and winter by simply turning off the basking lamp and letting room temperature prevail overnight. In summer or in consistently warm climates, a small fan directed at the exterior of the enclosure on a timer running at night can help the enclosure cool toward the lower end of that target. What you are working toward is an enclosure that cools meaningfully at lights-out and warms back up when the basking lamp switches on at sunrise. This daily thermal cycle contributes to appetite, normal reproductive cycling, shedding regularity, and general vigor in a way that a constant 75°F environment does not replicate.

UVB for an arboreal animal

Standard reptile UVB guides position the lamp over one end of a horizontal enclosure. For a tall arboreal chameleon enclosure, the primary UVB source goes at the top, aligned with the basking branch. The chameleon will spend its basking time 6 to 12 inches from the lamp surface — well within the correct UV index range for Ferguson Zone 3. Ensure the basking branch is positioned so the animal must actually choose to spend time near the lamp to receive it, as it would naturally choose to bask in sunlit vs. shaded branches.

Avoid the T5 HO 10.0 or ReptiSun 10.0 for veiled chameleons. That tube is appropriate for desert tortoises, large rock-dwelling agamas, and other Zone 4 species that bask in full unobstructed desert sun. A forest arboreal species living under partial canopy cover in highland mist — which is the veiled chameleon's actual natural context — does not receive Zone 4 UV intensity, and a chameleon basking six inches from a 10.0 tube is receiving a UV dose that has been linked to eye inflammation and corneal damage.

Replace the UVB tube every **6 months** without exception, and note the date on a label stuck to the enclosure. A lamp approaching the end of its useful UV life may look visually identical to a fresh lamp while delivering less than half the effective UV output.

SECTION 6

Feeding & Gut-Loading

A rotation beats a single staple: dubia roaches and crickets for the base, hornworms and silkworms for hydration and variety, black soldier fly larvae as a calcium-rich extra. Gut-load every feeder before it goes in.

Veiled chameleons are **strict insectivores** — with one important and biologically interesting exception: they consume live plant leaves. This is the most significant behavioral difference between Veileds and Panther Chameleons, and it has implications for both nutrition and plant selection. An otherwise well-fed veiled chameleon will browse leaves from pothos, hibiscus, and schefflera daily. This incidental plant consumption provides supplemental moisture and trace micronutrients, and it is one reason live enclosure plants are a functional part of the care setup rather than purely decorative.

The insect-based diet is non-negotiable. Unlike bearded dragons, which are genuinely omnivorous, a veiled chameleon supplemented with chopped vegetables as a primary food source will develop deficiencies and fail to thrive. The diet is insects, gut-loaded thoroughly.

Feeder insect guide

Feeder	Category	Notes
Dubia roaches	**Staple**	Excellent protein-to-fat ratio, easy to gut-load, low odor, no noise; the strongest general-purpose staple for chameleons of all ages

Feeder	Category	Notes
Crickets	**Staple**	High protein, widely available, stimulate active hunting behavior via movement and chirping; must be gut-loaded 24–48 hours; remove uneaten crickets within 30–45 minutes — nocturnal biting causes injury and stress
Black soldier fly larvae (BSFL, phoenix worms)	**Staple**	Naturally high calcium, soft-bodied, high moisture content supports hydration; excellent for juvenile and sub-adult chameleons
Hornworms	**Occasional treat**	Very high moisture, excellent hydration feeders, highly palatable; soft-bodied; grow large quickly — offer when small to medium-sized for most chameleons
Silkworms	**Occasional treat**	Excellent nutrition, very soft-bodied, high in protein; valuable for picky eaters, recovering animals, or as variety alongside staples
Superworms	**Occasional treat — adults only**	High fat and significant chitin content; too large for juveniles; a chameleon fed superworms regularly may begin refusing staple feeders. Use sparingly — no more than once a week maximum.
Waxworms	**Recovery use only — rarely**	Extremely high fat, highly palatable; useful for stimulating appetite in a convalescent or severely underweight animal; addictive. One to two maximum; not a component of normal rotation.
Mealworms	**Poor choice — minimal use**	Higher chitin content than dubias, lower nutritional density; occasionally acceptable as variety; not suitable as a primary or staple feeder
Wild-caught insects	**Never**	Pesticide and parasite vectors; captive-bred versions of the same species are strictly safer in every dimension
Chopped vegetables, fruit, or salad in a dish	**Never as a primary food offering**	Veiled chameleons may graze leaves from living enclosure plants but will not eat chopped greens from a bowl; do not attempt dietary supplementation this way

Gut-loading — the real nutrition delivery system

A feeder insect purchased and offered the same afternoon it arrives contains almost no nutritional value — its gut holds only shipping material or nothing at all. An insect that has been fed a high-quality diet for 24 to 48 hours passes that nutrition to the chameleon who eats it. This process is **gut-loading**, and it is not a refinement for advanced keepers — it is the mechanism by which your chameleon actually receives its vitamins, calcium, and trace minerals. Dust on the outside of a poorly-loaded insect is a thin nutritional layer over an empty shell.

Good gut-load items: mustard greens, collard greens, dandelion greens, squash, sweet potato, carrot, kale (in rotation, not exclusively), and a quality commercial dry gut-load blend available continuously. Avoid high-oxalate items like spinach and beet greens as the majority component. Replace fresh items daily; a dry commercial blend available at all times fills the gap.

Supplement schedule

Supplement	Frequency	Why
Calcium without D3	Every feeding — lightly dust feeders before offering	The most critical single supplement for captive chameleons; prevents metabolic bone disease (MBD) in animals that cannot synthesize enough calcium from insects alone
Calcium with D3	**2× per month**	D3 is fat-soluble and accumulates in tissue; twice-monthly dosing provides adequate levels for calcium absorption without the toxicity risk of more frequent supplementation
Reptile multivitamin	**1× per month**	Covers vitamin A, vitamin E, and trace minerals that gut-loading alone may not fully provide in all seasons and all feeder rotations

Do not over-supplement vitamin D3. Hypervitaminosis D3 — D3 toxicity — causes pathological soft tissue calcification and is a genuine and documented risk in reptiles whose keepers supplement more frequently than recommended. The twice-monthly schedule is deliberate: more D3 is not better.

Feeding schedule by age

Life stage	Age	Frequency	Notes
Hatchling	0–3 months	Daily	Small BSFL, pinhead-to-small crickets, or small dubia nymphs; feeder width must not exceed the distance between the eyes
Juvenile	3–6 months	Daily	Gradually increasing feeder size; introduce hornworms and silkworms for dietary variety
Sub-adult	6–12 months	Every 1–2 days	Build variety into the rotation; establish dubia roaches as the primary staple; observe hunting drive at each offering
Adult male	12 months+	Every 2 days	6–12 feeders per session; adjust down if body condition suggests excess fat around the casque or lateral folds
Adult female	12 months+	Every 2 days, adjusted for reproductive stage	Increase calcium supplementation frequency before and during visible egg development; watch closely
Gravid female	When visibly distended with developing eggs	Every 1–2 days with heavier calcium dusting	The highest-demand period for supplemental calcium; the egg-laying bin must be continuously available
Recovering or underweight	Any	Daily with softer-bodied feeders	Silkworms and hornworms preferred for ease of digestion; reassess after 30 days

Reading the chameleon during feeding

A healthy, interested veiled chameleon will lock both eyes forward onto a feeder insect, orient its body toward it, extend slowly in a stalking posture, and extend the tongue on a precise vector. A chameleon tracking feeders with its eyes but not committing to a strike may be slightly too cold (check the basking zone temperature), slightly dehydrated, or early-stage unwell. A chameleon who ignores feeders entirely — particularly one who hunted reliably the day before — needs an immediate temperature check, a

hydration assessment, and a vet consultation if no environmental explanation is identified within 24 hours.

Safe enclosure plants for incidental grazing

Since veiled chameleons actively consume enclosure plant leaves, every species in the enclosure must be confirmed non-toxic before introduction:

- **Pothos (*Epipremnum aureum*)** — the standard; very hardy under misting conditions
- **Hibiscus (*Hibiscus rosa-sinensis*)** — flowers and leaves both safe; widely accepted as beneficial
- **Schefflera / Umbrella plant (*Schefflera arboricola*)** — safe and widely used in chameleon setups
- **Ficus benjamina** — safe with thoroughly washed leaves; rinse the sap residue before introducing
- **Avoid:** Philodendron (toxic despite resembling pothos), any plant treated with pesticides or systemic fertilizers, dracaena species (toxic), dieffenbachia (toxic), all plants in the nightshade family, peace lily

> ***Shop the food and supplies:*** *Browse Reptile & Amphibian Food and Live Feed Delivered!.*

SECTION 7

Hydration — The Critical Section

Dehydration is the leading preventable cause of death in captive veiled chameleons. It is almost always the result of a keeper who did not fully understand how chameleons drink — not a keeper who was negligent, but one who applied the logic of every other reptile they had kept and found it did not translate. This section deserves its own heading.

How chameleons drink

Veiled chameleons in the wild drink from water droplets that accumulate on leaves, branches, and the surface of their own scales during morning mist events and rain. They drink by lapping individual droplets with the tongue — often for extended sessions, working systematically from leaf to leaf, collecting dozens of small water contributions over several minutes. The casque on the head channels water toward the mouth during these drinking sessions, a functional anatomy adaptation to the mist-heavy highland environment.

Many captive chameleons will not recognize standing water in a dish as a drinking source at all. Some individuals can be trained to drink from a drip system aimed at a leaf or the side of the enclosure, but no chameleon should be expected to find its primary hydration in a bowl.

Your hydration delivery system is the misting schedule, and it must work correctly. Misting must coat leaf surfaces throughout the enclosure — not only the walls — for long enough that the chameleon can drink. A minimum of two automated misting cycles daily (morning and evening), each running 3 to 5 minutes, is required for most setups. In dry rooms, summer months, or warm climates, a midday session may also be needed.

A **drip system** running for 1 to 2 hours daily between misting cycles supplements the automated schedule. A dripper landing on a leaf at mid-enclosure height provides a continuous drinking opportunity for a chameleon that is reluctant to drink during or immediately after the burst misting cycle. This is particularly valuable during the first weeks with a new animal that is still learning the enclosure's routines.

Signs of adequate hydration

- Urates (the white portion of droppings) are white or very pale yellow
- Eyes are round, full, and flush with the turret margins — not recessed
- Skin is smooth and returns immediately when very gently displaced
- The chameleon is alert, holds its position on branches confidently, and browses the enclosure
- Drinking behavior is observed during or after misting — tongue-lapping at leaf surfaces

Signs of dehydration — act immediately

- **Sunken eyes** — the most reliable early indicator; the eye turrets appear to have recessed backward and inward; the eyes look smaller and the area around the turret base appears hollow. This is an urgent concern, especially when combined with any other symptom.
- **Orange or dark yellow urates** — signals the chameleon is not taking in enough water relative to metabolic demand
- **Lethargy combined with loss of bright coloration** — a dark, inactive chameleon that was green and active yesterday
- **Skin tenting** — gently displace a small fold of skin on the lateral body; in an adequately hydrated animal it returns to position immediately; in a dehydrated animal it holds the tented shape for a brief but visible moment before returning
- **Sticky or ropy mucus** visible as threads inside the mouth when the animal opens it

What to do for suspected dehydration: increase misting immediately — extend session lengths and add a midday cycle. Verify with your own eyes that misting nozzles are wetting leaf surfaces throughout the enclosure, not only coating the screen walls. A chameleon showing sunken eyes should be seen by a vet within 24 hours if symptoms do not begin resolving with corrected misting; oral fluid supplementation by a vet can be life-saving in moderate to severe dehydration.

The warm-mist shower as a supplemental tool

Some experienced keepers periodically place a chameleon on a branch or plant in a bathroom and allow warm steam from a running shower — not the direct water stream — to fill the room for 10 to 15 minutes. Many chameleons respond by drinking extensively from the condensation that collects on branches and their own body surface. This is a useful supplemental technique for an animal that seems chronically under-drinking, but it is not a substitute for a correctly designed in-enclosure misting system — it supplements, not replaces.

SECTION 8

Shedding

Adult veiled chameleons shed every **4 to 8 weeks**, a cycle that compresses to every **2 to 4 weeks** in rapidly growing juveniles. Before a shed, the coloration becomes notably dull, slightly gray or pale, and the chameleon may reduce activity and appetite for a day or two. Both of these changes are normal pre-shed behavior, not a health concern.

What normal shedding looks like

Chameleons shed in sections rather than in a single intact skin the way snakes do. The shed lifts and peels from the casque and head first, then works its way down the body, and finishes at the feet and tail tip. Most of the process happens over 24 to 48 hours, often concentrated overnight. Unlike leopard geckos, veiled chameleons do not typically consume their shed skin.

Increase misting frequency in the **24 to 48 hours leading up to and during a shed**. Additional surface moisture helps the shed lift cleanly away from the new skin beneath. Dry conditions during shed translate directly into stuck shed. A chameleon in active pre-shed should not be handled — the loosening skin is sensitive and disrupting the shed cycle adds unnecessary stress.

Stuck shed and dysecdysis

Dysecdysis (dis-EK-di-sis) is the term for incomplete or retained shed. In chameleons, the two most consequential sites are the **turret eyes** and the **toe tips**.

Retained eye caps: each eye turret has a specialized shed layer that must release cleanly with the rest of the head shed. A cap that remains adhered over a turret eye constricts the eye, impairs vision, and can cause permanent damage or secondary infection if left in place. It presents as a dull, non-responsive, slightly recessed-looking eye area with a visible dried layer. Do not attempt to peel it away manually. Increased misting aimed directly at the face from a comfortable distance may help loosen it over 24 to 48 hours. If the eye cap has not released on its own by then, this is a vet appointment.

Retained shed on feet and toes: the zygodactyl feet have numerous skin folds around the toe bundles where old shed accumulates across multiple shed cycles. Check all four feet carefully after every shedding event. Constricting rings of retained shed cause progressive circulatory restriction and ultimately toe loss — a permanent outcome that is entirely preventable with consistent post-shed checks. Extended gentle misting and a warm damp cloth held loosely around the affected foot for several minutes can loosen retained material. If constriction is already visible as a visible ring or pale distal toe segment, see a vet promptly.

What not to do: do not attempt to peel dry shed from any area of the chameleon's body. Dry shed is adhered to new skin. Pulling it before it has been sufficiently moistened tears the new skin, exposes raw tissue to infection, and causes significant discomfort. Always moisten the area thoroughly first, allow several minutes for it to soften, and attempt only gentle loosening with a damp cotton swab. If the shed does not release easily, stop, mist more, and try again the following day.

SECTION 9

Common Challenges & Solutions

Veiled chameleons are demanding precisely because environmental errors compound quickly and illness hides effectively. The problems below are the ones every keeper needs to know before they arise.

Dehydration — sunken eyes

What you see: the eye turrets appear recessed or sunken into the head; the chameleon is lethargic, slow to react, and coloration is dull; urates are orange or yellow rather than white. **Why:** the chameleon is not taking in enough water — most commonly because misting cycles are too short, misting hits only walls rather than leaf surfaces, no drip system is running, or the animal is newly arrived and still learning where to find water in this enclosure. **What to do:** increase misting frequency and duration immediately; confirm with your own eyes that droplets are landing on branches and leaf surfaces throughout the enclosure; start or extend a drip system; assess urate color at every observation. **When to call a vet:** any chameleon showing sunken eyes who does not show improvement within 12 to 24 hours of corrected misting — oral fluid supplementation by a vet can reverse moderate dehydration rapidly.

Metabolic bone disease (MBD)

What you see: trembling or shaking limbs during movement; a lower jaw that appears soft or rubbery when viewed; bowing of the legs or spine; difficulty gripping branches; in severe cases, spontaneous fractures or an animal that can no longer support its own weight. **Why:** calcium or vitamin D3 deficiency — most commonly from inadequate calcium dusting at feedings, a failed or aged-out UVB tube, the UVB tube being too far from the basking zone to deliver Zone 3 UV, or a combination. Females during and after clutch production are at significantly elevated MBD risk because the calcium demand of egg shell formation is substantial. **What to do:** correct the supplement schedule immediately; verify the UVB tube is within its first 6-month window; confirm the basking branch is 6 to 12 inches from the tube. **When to call a vet:** as soon as trembling limbs or jaw softness are visible — MBD responds well to early intervention and is very difficult to reverse once bone deformity is established.

Respiratory infection

What you see: mucus visible at the mouth corners or nostrils; the chameleon periodically gapes (holds its mouth open while resting on a branch); audible clicking or wheezing from the chest during breathing; lethargy; reduced or absent basking even when temperatures are correct. **Why:** bacterial or viral respiratory infection, almost always precipitated by a glass or blocked-screen enclosure, a misting cycle that never fully dries between sessions, stagnant high humidity without air movement, or a consistently cool ambient temperature. **What to do:** improve enclosure ventilation immediately; ensure the wet-dry humidity cycling is functioning as designed; correct any temperature drift; see a vet promptly. **When to call a vet:** same day or next day — respiratory infections in chameleons escalate rapidly and almost always require antibiotic treatment to resolve.

Egg-binding (dystocia) in females

What you see: a visibly distended female who has been digging persistently in the enclosure floor or attempting to dig against the walls for more than 24 to 48 hours without depositing a complete clutch; she appears exhausted or lethargic; straining efforts are visible. **Why:** eggs cannot be passed because no suitable laying site was provided, because an egg is malpositioned in the reproductive tract, or because calcium depletion from previous clutches has weakened the uterine muscle contractions needed for laying. Veiled chameleons produce infertile clutches without any male present — this is biologically normal and expected, which means a deep laying bin must be a permanent fixture for any adult female. **What to do:** ensure the laying bin is continuously in place — a container at least 12 inches deep, filled with moist 60:40 soil-and-sand mixture, placed at the enclosure floor level with an entry hole. A female preparing to lay needs privacy, warmth, and adequate digging depth. Warm soaks may assist in very early cases. **When to call a vet:** any female who has been actively digging or straining for more than 48 hours without completing a clutch — dystocia in chameleons is a veterinary emergency.

Internal parasites

What you see: chronic weight loss despite apparently normal appetite; loose, malformed, or off-color droppings over multiple weeks; a generally unthrifty appearance and dull coloration that does not improve with environmental corrections. **Why:** internal parasites — pinworms, flagellates, coccidia — are common in chameleons acquired through commercial channels, reptile expos, or any facility where hygiene cannot be individually verified. Newly imported or recently wild-caught animals are at higher risk. **What to do:** take a fresh fecal sample to an exotic vet for microscopic examination — make this a standard appointment within the first 30 days of ownership for any new chameleon, regardless of apparent health. **When to call a vet:** baseline fecal exam for every new animal; for the above symptoms in any established animal, within a week.

Thermal burns

What you see: blistered, lifted, or discolored scales along the dorsal ridge, casque surface, or upper body; the chameleon may be spending unusual amounts of time lower in the enclosure; affected scale areas may look gray, yellow-tinged, or in severe cases appear as raised blisters or necrotic patches. **Why:** the basking branch surface temperature exceeded 95°F, or the basking lamp was positioned too close to the top screen, creating an overly intense focal hotspot on the primary basking branch. Chameleons basking directly under the light can be slow to detect developing skin damage and may stay in a burning position longer than a mammal would. **What to do:** measure the basking branch surface immediately with an infrared gun; if reading above 95°F, reduce bulb wattage or increase the lamp-to-branch distance; move the chameleon to a clean, correctly-temped enclosure while the affected skin heals. **When to call a vet:** any burn showing lifted scale edges, tissue discoloration extending beneath the scale surface, or any burn involving the casque or eyes — burns in reptiles become infected rapidly and often require antibiotic support.

Stomatitis (mouth rot)

What you see: the chameleon is repeatedly wiping its mouth on branches or screen panels; gum tissue around the mouth is reddened or visibly swollen; a yellowish or cheesy-looking discharge is present at the lip margins; in more advanced cases, blackened or necrotic tissue visible at the gum line. **Why:** bacterial infection initiated by a small wound from a cricket bite overnight, abrasion from rough enclosure surfaces, or the immune suppression that accompanies chronic stress or poor environmental management. **What to do:** examine the mouth in good lighting; gently clean accessible areas with a dilute chlorhexidine solution (1:10 dilution) on a cotton swab; eliminate overnight live feeders that may bite; address any chronic environmental stressors. **When to call a vet:** within two to three days of noticing first symptoms — stomatitis spreads into jaw bone rapidly in chameleons and becomes life-threatening without antibiotic intervention.

Chronic stress coloration — dark all day

What you see: a chameleon that is persistently dark brown to near-black through most of the day, rarely or never showing the bright greens and yellows of a settled, contented animal; may also hiss, gape defensively, or rock side-to-side when the keeper approaches; may stop eating within a few days of onset. **Why:** something in the environment is causing persistent unresolvable stress — most commonly visual exposure to another chameleon through a shared room, the chameleon's own reflection in a nearby window or reflective surface, a household dog or cat visible to the animal, an enclosure in a high-traffic corridor, too many handling attempts too early, or an environmental parameter that has been consistently out of range. **What to do:** systematically audit potential stressors; check all temperatures and humidity cycling; add visual coverage inside the enclosure; assess the room position and adjacent animal traffic; reduce handling to zero during the investigation. **When to call a vet:** a chameleon who stays dark, refuses food, and shows no improvement 3 to 5 days after environmental stressors have been corrected — chronic stress causes immune suppression and opens the door to secondary infection.

Visual stress from external sources

What you see: similar to chronic stress coloration but often episodic or time-correlated — dark coloration and defensive posturing that triggers at specific times of day, when certain people or animals enter the room, or in consistent lighting conditions. **Why:** the chameleon's 340-degree vision is detecting something outside the enclosure that it registers as a threat — another chameleon in the room, a dog or cat passing by, outdoor birds visible through a window, or a reflection. Because the perceived

threat never resolves (it keeps returning), the chameleon cannot complete a normal defensive cycle and remains on elevated alert. **What to do:** identify the trigger through timing observation; block the relevant line of sight with a solid background panel, an enclosure repositioning, or restricted pet access to the room. **When to call a vet:** not typically a vet issue on its own if resolved promptly; a chameleon who develops secondary health changes from prolonged stress-related immune suppression should be evaluated.

Quick reference

What you see	Likely cause	What to do	Vet urgency
Sunken eyes, orange urates	Dehydration	Increase misting immediately; add drip system	12–24 hours if no improvement
Trembling limbs, soft jaw	MBD	Correct supplements and UVB immediately	Within days; sooner if severe
Mucus at mouth or nose, gaping	Respiratory infection	Improve ventilation; correct temperatures	Same day or next day
Female digging 48+ hours without laying	Egg-binding / dystocia	Provide deep laying bin; warm soak	Same day if straining continues
Weight loss with loose droppings	Internal parasites	Fecal exam	Within a week; sooner if declining
Blistered or lifted dorsal scales	Thermal burn	Measure and correct basking temperature	Tissue lifting or necrosis: same day
Cheesy discharge at mouth margins	Stomatitis	Chlorhexidine clean; eliminate biting feeders	Within 2–3 days
Dark coloration all day, not eating	Chronic environmental stress	Audit systematically; reduce handling	3–5 days if no resolution after fixes
Dark coloration at consistent times	Visual stress from external source	Identify and block line-of-sight trigger	Monitor; vet if secondary illness develops

SECTION 10

Handling

The most important thing to understand about handling a veiled chameleon is this: **they tolerate it, they do not enjoy it.** A well-habituated adult veiled chameleon may come to accept regular brief handling with minimal visible stress — and that is the realistic goal. Not a chameleon that seeks contact with you, but a chameleon that is not in distress during it. Setting this expectation clearly from the beginning prevents a great deal of misreading the animal's behavior and a great deal of chronic stress caused by well-meaning keepers who pushed for more interaction than the animal could give.

The minimal handling philosophy

Chameleons communicate stress clearly and consistently, if you know the signals. A chameleon gaping, turning dark, rocking side-to-side, or hissing is asking to be put back. Continuing to hold a chameleon past the point of visible stress does not produce a tamer animal over time — it produces a chameleon with elevated baseline cortisol and a weakened immune system. Shorter sessions that end when the animal signals it has had enough produce better long-term results than longer sessions that end when the keeper decides.

Aim for **5 to 10 minutes per handling session** maximum, particularly during the first three months. Handle no more than every two to three days during the habituation period. A chameleon who is eating reliably, basking confidently, and showing good color is a chameleon who should not be handled more simply because it will tolerate it — routine observation from outside the enclosure is sufficient for a healthy, settled animal.

How to pick up a chameleon

Never reach down from above toward a chameleon. The aerial approach mimics a bird of prey and triggers an immediate defensive response — it is the most reliable way to produce a gaping, dark, hissing animal. Instead, approach at the chameleon's level from the side, present your hand open and palm-up below the branch it is on, and gently raise your hand toward the branch. A chameleon that steps onto your hand voluntarily is indicating it accepts the interaction. A chameleon that has to be peeled off a branch is telling you clearly that it does not.

Support the full body. Allow the chameleon to walk from hand to hand at will. Handle seated and low — over a couch, over your lap — never at standing height. Chameleons can fall and do fall, and a drop of two feet onto a hard floor can cause serious injury. Bring the chameleon back to the enclosure at the first stress signal; do not wait for the second one.

Stress signals

- Dark coloration — particularly if the animal was green when you opened the enclosure
- Gaping (holding the mouth open while at rest — not a yawn, a threat display and a respiratory signal)
- Hissing or audible breath
- Rocking the body side-to-side (a defensive behavior mimicking a leaf in the wind)
- Flattening the body laterally to appear as thin as possible (a threat display)
- Persistent movement in one direction away from you

Any of these signals means the session ends now. Return the chameleon to the enclosure. Do not resume the session the same day.

Children and veiled chameleons

Veiled chameleons are not well-suited as pets where young children are the primary caregivers. They are visually captivating, slow-moving, and undeniably fascinating to observe — but they require from the handler a stillness, patience, and sensitivity to subtle behavioral signals that most children under ten do not reliably have. A chameleon grabbed, held too long, or carried into a

high-stimulus environment will experience real physiological stress. For families with children, close observation of a well-planted enclosure is genuinely enriching — arguably more so than direct handling — and carries none of the stress risk. If handling does occur with children present, the child should be seated on the floor, remain very still, and return the animal the moment any stress signal appears.

SECTION 11

Routine Care Calendar

How often	What to do
Daily	Run the misting system morning and evening; confirm with direct observation that misting is coating leaf surfaces, not only walls
Daily	Check the basking spot temperature with an infrared gun; confirm the UVB lamp is cycling on and off on the correct photoperiod
Daily	Observe the chameleon for 5 minutes: coloration, activity level, eye fullness, any mucus at mouth or nostrils, posture on branch
Daily	Offer food on the appropriate schedule; remove any uneaten live insects from the enclosure within 30–45 minutes
Daily	Check urate color in droppings — white is normal; orange or yellow flags inadequate hydration
Daily	Spot-clean the drainage tray or floor area; remove dead plant matter and waste accumulation
Daily	Confirm cool end temperature with a probe thermometer
Weekly	Full inspection of all four feet and all toe tips for retained shed, swelling, or constriction
Weekly	Close visual check of the casque and eye turrets — any swelling, discoloration, or retained shed around the turret margins
Weekly	Clean misting nozzle heads; mineral deposits gradually restrict misting output and are the most common cause of misting system failure
Weekly	Replace drip reservoir water with fresh filtered or dechlorinated water; rinse the reservoir
Weekly	Check feeder insect supply; reorder before running low; confirm gut-load is fresh
Weekly	Audit supplement rotation — confirm calcium, calcium+D3, and multivitamin are cycling correctly
Weekly	Assess live plant health — trim dead or yellowing leaves; check for root rot at pot drainage holes
Monthly	Deep clean — remove all branches and decor; wipe screen panels with reptile-safe disinfectant; rinse and allow to dry fully
Monthly	Repot or rotate live plants if root-bound or showing nutrient stress from misting water mineral buildup
Monthly	Wash all decor, branches, cups, and vines with reptile-safe disinfectant; rinse thoroughly and dry before returning
Monthly	Photograph the chameleon in lateral view for a body condition baseline to compare over time
Monthly	Review supplement supply and reorder before running low
Every 6 months	Replace the UVB tube — mark the installation date on a label stuck to the enclosure or in a log
Annually	Exotics vet wellness check; fecal exam; body weight record; review supplement schedule and husbandry with the vet

SECTION 12

When to See a Vet

Chameleons mask illness as a survival behavior inherited from prey animals. By the time visible symptoms appear, the underlying problem has almost always been developing for a meaningful period. The rule is simple: **when you are not certain whether to call, call.** A five-minute phone consultation costs nothing compared to a chameleon who crossed a critical threshold while you were deciding.

Go today — same day or emergency:

- Sunken eyes combined with lethargy and absence of normal color
- Open-mouth breathing or gaping at rest; audible clicking or wheezing from the chest
- Active straining in a female who has been digging for more than 24 to 48 hours without completing a clutch
- Any tissue prolapsing from the cloaca (rear vent)
- A chameleon found on the enclosure floor, unable to climb, or unable to right itself on a branch
- Sudden inability to extend or retract the tongue
- An obvious fracture, wound with exposed tissue, or persistent bleeding
- Thermal burn showing lifted scales, tissue discoloration, or necrosis
- Sudden complete coordination loss, circling, or seizure-like activity

Book within a few days:

- Urates that are consistently orange or yellow despite 24 hours of corrected and verified misting
- Retained eye shed that has not released after 48 hours of increased misting and humidity
- A lump or unexplained swelling anywhere on the body
- Mouth tissue that is red, swollen, or showing any discharge
- Any progressive visible weight loss — a chameleon who has become noticeably thinner over two to three weeks
- Loose or malformed stool over multiple days combined with reduced or absent feeding drive
- A chameleon who has stopped hunting entirely for more than 4 to 5 days with no identifiable environmental cause

Find your exotic animal vet before your chameleon needs one. When you call, confirm specifically that the clinic has experience with chameleons — not just reptiles generally. Bring a written temperature log, your misting schedule, your supplement rotation record, and a fresh fecal sample in a sealed container to every appointment.

SECTION 13

FAQ

Are veiled chameleons good pets for beginners?

Veiled chameleons are the most beginner-accessible species within the chameleon family — more forgiving than a Jacksons, a Mellers, or a Fischer's — which makes them a reasonable starting point for someone who has specifically decided to keep a chameleon. But that framing is important: they are the entry point for chameleon keeping, not the entry point for reptile keeping. Their ventilation, hydration, temperature, and UVB requirements are all precise, and they conceal illness until it becomes critical. Someone who has kept a forgiving reptile like a bearded dragon or leopard gecko for at least a year, has done serious advance research, and is prepared to invest in a proper screen enclosure and an automated misting system will find veiled chameleons genuinely rewarding. Someone who is new to reptiles and looking for a low-maintenance companion should start elsewhere and return to chameleons when they have a foundation to build on.

What do chameleons eat?

Veiled chameleons are strict insectivores, with the interesting exception that they will also graze live plant leaves from vegetation growing inside their enclosure. The core diet is insects, all of which should be gut-loaded on fresh leafy greens and a quality dry gut-load blend for 24 to 48 hours before offering. The best staple feeders are dubia roaches, crickets, and black soldier fly larvae. Hornworms and silkworms make excellent occasional additions for variety and hydration. Calcium powder should be dusted onto feeders at every meal, with calcium-plus-D3 used twice a month and a multivitamin once a month. Do not offer chopped vegetables, fruit salad, or greens in a dish — veiled chameleons eat plant material only by grazing living enclosure plants directly, not from a bowl.

Do chameleons need a screen cage?

Yes — without exception, and the reason is biological rather than preferential. Glass terrariums trap heat, stagnant air, and moisture, and they prevent the wet-dry humidity cycling that veiled chameleons require. In a glass enclosure with a running mister, the warm, stagnant, persistently humid microenvironment becomes a reliable culture medium for the bacteria that cause respiratory infections. Chameleons have thin respiratory passages and are highly sensitive to poor air quality. The minimum enclosure for one adult is 24 inches wide by 24 inches deep by 48 inches tall — all-screen or mesh construction on multiple faces.

How do you give a chameleon water?

Chameleons drink from water droplets on leaves and surfaces — not from standing bowls. The primary delivery method is an automated misting system programmed to run at least twice daily, morning and evening, with each session lasting long enough to coat leaf surfaces throughout the enclosure with visible droplets. Most keepers supplement with a slow drip system running for one to two hours daily between misting sessions, positioned to drip onto a leaf at mid-enclosure height. Some keepers also use a warm-steam shower environment periodically — allowing the chameleon to sit near a running shower in a misted bathroom for 10 to 15 minutes — which encourages many animals to drink extensively. Standing water dishes are not harmful to include but cannot be relied on as the sole water source.

What temperature do chameleons need?

The basking branch surface directly under the heat lamp should sit at **85–95°F** — measure it with an infrared temperature gun. The ambient warm-side air should be **75–80°F** and the cool end **68–72°F**. At night, when the basking lamp switches off, the enclosure should drop to **55–65°F**. This overnight temperature drop is one of the most surprising requirements for first-time chameleon keepers and one of the most important ones for long-term health — veiled chameleons are highland animals that evolved under significant day-to-night temperature swings, and a room that stays at 75°F all night disrupts their circadian rhythm and long-term metabolic function. Every heat source must be on a thermostat; an unregulated basking lamp can push the branch

surface above 95°F and cause burns.

How long do veiled chameleons live?

Male veiled chameleons typically live **5 to 8 years** with good husbandry. Females live a shorter **3 to 5 years** because egg production — which occurs even in females that have never been near a male, as veiled chameleons routinely produce infertile clutches — places significant ongoing demand on calcium stores and body reserves. These lifespans are shorter than most reptile keepers expect, especially compared to tortoises or leopard geckos, and it is worth factoring that in realistically before acquisition. The most reliable predictors of a lifespan toward the upper end of those ranges are correct environmental management from day one, consistent calcium supplementation particularly in females, avoidance of over-breeding females, and annual veterinary wellness checks.

Why is my chameleon dark colored?

Color in veiled chameleons communicates thermoregulation status, mood, and health — it is not exclusively about camouflage. A chameleon that is dark brown first thing in the morning while basking is absorbing heat and behaving normally. A chameleon that stays dark through its active period, remains dark at rest in a correctly heated enclosure, or is dark and lethargic together is communicating that something is wrong. The most common causes of persistent dark coloration are chronic stress from visual exposure to another animal or its own reflection, an enclosure position near high household traffic, environmental parameters that are consistently out of range, or early-stage respiratory or systemic illness. A chameleon that stays persistently dark for more than one to two days after obvious environmental causes have been ruled out and corrected should be seen by a vet.

Do chameleons need UVB light?

Yes — veiled chameleons are Ferguson Zone 3 reptiles with high UVB requirements. In the wild they bask in direct highland sunlight for extended periods each day, receiving UV indices well above what a shaded or indoor environment delivers. A T5 HO 5.0 or 6% fluorescent tube positioned so the basking branch is 6 to 12 inches from the lamp surface delivers the correct UV index. Avoid the 10.0 or 12% tubes — these deliver a UV index appropriate for desert species, not forest arboreal animals, and at close basking distances they have been linked to corneal and eye inflammation. The UVB tube must be replaced every 6 months regardless of whether it appears to be lighting normally, because effective UV output degrades before visible light output does.

Can two chameleons live together?

No. Veiled chameleons are strictly solitary, and this applies not only to physical cohabitation but to visual contact. Two chameleons that can see each other through adjacent enclosure panels will both experience persistent territorial stress — the threat is always there and never resolves, which means the stress response never completes. This causes chronic immune suppression, reduced feeding drive, and shortened lifespan in both animals. Every chameleon requires its own enclosure placed where it cannot see any other chameleon in the room. There are no practical exceptions — not two females, not juveniles, not a male and female except for brief supervised breeding introductions, and not two enclosures simply placed a few feet apart.

How often should I mist my chameleon?

At minimum, twice daily — once in the morning and once in the evening — with each session running long enough (3 to 5 minutes) to thoroughly wet leaf surfaces throughout the enclosure, not only the screen panels. In dry climates, rooms with low ambient humidity, or during summer months, a midday misting session may also be required. The key indicator is urate color: white urates mean the chameleon is adequately hydrated; orange or yellow urates mean misting needs to increase immediately. A supplemental drip system running for one to two hours daily between misting cycles significantly improves outcomes, especially for animals that are slow or reluctant drinkers. Watch for tongue-lapping behavior at leaf surfaces during and after misting — that is confirmation that the system is working and that the chameleon is drinking from it.

Ready to shop? *Start with Reptiles & Amphibians. 10% of every purchase supports no-kill animal shelters. Not sure this is the right pet? Compare your options in our Reptile Guide.*

SECTION 14

Every Purchase Gives Back

Before we earn a dime, **10% of every purchase at Walnut Creek Supply goes directly to no-kill animal shelters across the country.** Not at checkout. Not as an optional add-on. It comes off the top of every single order, every single day — because we believe that anyone committed enough to read a full chameleon care guide is someone who takes every animal seriously, not just the one at home.

We are a family in Christiana, Lancaster County, Pennsylvania. We grow our own food, keep our own animals, and stock the clean-label, American-made goods we would put in front of our own pets — **no mystery ingredients**, no glass tank marketed as a chameleon setup, and no 10.0 UVB tube sold without an honest explanation of why the 6% is the right call. If you are wrestling with a misting schedule, trying to read your chameleon's coloration, or want a second opinion on a supplement rotation, call us at **(717) 983-8069**. We would rather talk you through it.

Veiled chameleons reward the keepers who get the environment right and respect the animal's nature. There is nothing quite like a settled male firing up brilliant greens and yellows on his basking branch on a clear morning, completely at ease in his highland-mountain-in-a-box, hunting a dubia with that locked-on binocular stare, while the misting system finishes its morning cycle and the leaves drip. That moment is yours to earn — and this guide is how you get there.

From our family to yours, **Walnut Creek Supply** — Christiana, Pennsylvania *walnutcreeksupply.com*