

WALNUT CREEK SUPPLY

THE COMPLETE CARE GUIDE

Corn Snake

The ideal first snake — setup, feeding and long-term care made simple



Family-owned in Christiana, Lancaster County, Pennsylvania

10% of every purchase supports no-kill animal shelters nationwide

13 sections · walnutcreeksupply.com

CONTENTS

What's Inside

-
- 01 **Quick Start: Your First 48 Hours**
-
- 02 **Meet Your Corn Snake**
-
- 03 **Enclosure Setup & Layout**
-
- 04 **Substrate Guide**
-
- 05 **Heat, UVB & Humidity — The 3 Numbers**
-
- 06 **Feeding & Nutrition**
-
- 07 **Shedding**
-
- 08 **Common Challenges & Solutions**
-
- 09 **Handling**
-
- 10 **Routine Care Calendar**
-
- 11 **When to See a Vet**
-
- 12 **FAQ**
-
- 13 **Every Purchase Gives Back**

New owner? Go straight to Section 1. It is written so you can set up a safe, correct home and get through the first 48 hours without reading anything else first.

*Corn snakes are quietly the best beginner snake in the hobby — not the most visually dramatic on an expo table, not the most talked-about on reptile forums, but consistently the one that actually works. An adult corn snake in a well-furnished enclosure is active, curious, and genuinely exploratory in a way that makes keeping her engaging from day one. She investigates every corner of the setup, accepts food on schedule with far less of the temperamental, months-long fasting cycles that define ball python ownership, and emerges from her hide readily when the enclosure opens. She is slender, fast, and endlessly interested in her environment. The catch is that "easiest to keep" is sometimes read as "forgiving of corners cut," and that is where problems start. Corn snakes are extreme escape artists — every lid must latch securely, and gaps that look impossible are gaps they find with methodical patience. They require a real thermal gradient, a humidity range that prevents **dysecdysis** (incomplete shedding, defined in the Shedding section), and strictly sized frozen-thawed prey throughout their life. None of this is complicated. It simply needs to be correct from day one and maintained without drift. You may have this snake for fifteen to twenty years. Start right, and she will spend most of that time giving you very little to worry about.*

SECTION 1

Quick Start: Your First 48 Hours

1. **Build the complete enclosure before the snake arrives.** Minimum footprint for a juvenile or adult is 4' × 2' × 2'. Corn snakes are active and exploratory — unlike ball pythons, they use floor space enthusiastically. A juvenile in a correctly furnished 4-foot enclosure does fine; if feeding refusals occur, reduce visible open floor area with additional hides and foliage rather than downsizing the enclosure.
2. **Establish the thermal gradient and verify it with probe thermometers before any snake goes in.** Warm side surface temperature **85–90°F at substrate level**. Warm ambient air **80–85°F**. Cool end **72–78°F**. The gradient should span the full length of the enclosure — it is a spectrum, not two isolated zones.
3. **Connect every heat source to a thermostat — no exceptions.** An overhead halogen bulb needs a dimmer thermostat. A radiant heat panel needs a dimmer thermostat. An under-tank mat used as secondary heat needs an on/off thermostat with its probe placed between the mat and the enclosure floor. **Never use a heat rock.** Heat rocks develop unpredictable hot spots and have no safe application in any snake enclosure at any life stage.
4. **Set enclosure humidity to 40–60% before the snake arrives.** Corn snakes need moderate humidity — significantly lower than ball pythons. Aspen shavings, coconut fiber, or cypress mulch at 2–3 inches depth holds this range in a reasonably well-sealed enclosure without additional misting under most household conditions. Place a digital hygrometer at mid-enclosure and let the setup stabilize for 12–24 hours. Plan to raise humidity to 60–70% during shed cycles.
5. **Add two snug-fitting opaque hides before the snake arrives.** One on the warm side, positioned over the warmest part of the substrate. One on the cool end, in the 72–78°F zone. Each hide should require the snake to slide in and feel wall contact on most sides — corn snakes feel secure through tight confinement, not open shelter. A hide three sizes too large provides no security.
6. **Fill the water dish and place it on the cool side.** Corn snakes drink readily and will soak occasionally, particularly before a shed. Use a stable ceramic or stone dish that cannot be overturned. Refresh daily; wash thoroughly once a week.
7. **Secure every lid latch before the snake goes in.** Test the full seal by pressing gently on each section. Corn snakes are legendary escape artists. They detect temperature differentials through gaps barely wider than their head and their slender build allows them to push through spaces that look impossible. A screen lid needs clips on every side. A glass or PVC enclosure needs every door lock fully engaged. Check it every single time you close the enclosure.

8. **Place the snake in the enclosure and leave her alone for five to seven full days.** No handling, no feeding attempts, no rearranging the furniture. She needs to locate her hides, map the gradient, and establish that the space is safe. Check water and temperatures quietly once a day. Everything else can wait.
9. **Offer the first meal on day seven or later.** Use a frozen-thawed or freshly pre-killed rodent sized to the snake's mid-body girth. If she refuses, wait five to seven days and try again. Some corn snakes take two to three weeks to eat in a new home. Most settle and feed more quickly than ball pythons, but patience still applies.
10. **Find an exotic reptile vet before you need one.** Confirm they see colubrids specifically — not every exotic practice has experience with corn snakes. Keep the number somewhere accessible. A corn snake who looks mildly unwell on Monday can be in genuine trouble by Thursday.

First-Time Starter Bundle

What needs to be in the house on day one, and what to look for in each category.

Essentials

- **An enclosure with the correct footprint** — 4' × 2' × 2' for one adult, with a lid or door that seals completely and latches securely. PVC reptile enclosures and glass terrariums both work at this humidity range; glass requires less humidity management at 40–60% than it does for a ball python at 60–80%. The lid security is the non-negotiable element, not the material.
- **An overhead heat source on a thermostat** — a halogen flood lamp, ceramic heat emitter (CHE), or radiant heat panel wired to the appropriate thermostat type. The thermostat probe goes on the warm-side floor surface where the snake rests. Never power any overhead heat source without a thermostat in the circuit.
- **Two calibrated digital probe thermometers** — one at warm-side substrate level, one at cool-end floor level. An infrared temperature gun is useful for spot-checking surface temperatures during the setup phase. Stick-on thermometer strips are not a substitute for probe readings.
- **A digital hygrometer** — positioned at mid-enclosure, away from the water dish.
- **Two opaque hides** — one warm-side, one cool-side. Fully enclosed on all sides except a single entrance. Snug-fitting is the specification: a corn snake is not comfortable in a cave that feels open.
- **Substrate at 2–3 inches depth** — aspen shavings, coconut fiber, or cypress mulch. Cedar and pine contain aromatic phenols toxic to reptiles and must never be used at any life stage.
- **A stable water dish** — ceramic or stone, cool-side, wide enough for the snake to partially soak before a shed, stable enough to resist the snake's weight during movement.
- **Secure lid clips or door locks on every panel** — the single most important item on this list and the one most consistently skipped.

Ready to set up? Browse the essentials in our collections: [Reptile and Amphibian Enclosures and Reptile and Amphibian Terrarium Supplies & Accessories](#).

Enrichment

- **Additional hides and shelters** — corn snakes use floor space actively and benefit from a third hide or multiple covered resting spots across the enclosure. More cover options produce a calmer snake and a more confident feeder.
- **Cork bark pieces and tubes** — flat cork bark provides a natural resting surface with enough texture to help initiate a shed; cork tubes offer an enclosed elevated perching point the snake will reliably use.

- **Climbing branches and driftwood** — securely anchored diagonal branches at varying heights. Corn snakes are legitimate climbers and will use elevated perches routinely during their evening active period. Any branch must be fixed so it cannot shift under the snake's full body weight moving at speed — a fall from an unstable branch causes spinal injuries.
- **Dense artificial foliage** — placed along the back and sides to break open sightlines, reduce ambient stress, and give the snake a sense of visual cover while moving through the space.
- **A shallow sphagnum moss zone** — a small covered area packed with lightly damp sphagnum moss, available during shed cycles. Keep it contained so the rest of the enclosure maintains its normal 40–60% baseline.
- **A ventilated travel carrier** — a snug-fitting plastic container with a secure lid and a small hide inside. Introduce it to the home enclosure weeks before any vet trip so the carrier is a familiar scent rather than an alarm signal.

Enrichment, one click away. Shop Reptile and Amphibian Accessories.

SECTION 2

Meet Your Corn Snake

Wild corn snakes (*Pantherophis guttatus*) range across the eastern and central United States — from New Jersey south through Florida and west through Tennessee, Oklahoma, and Louisiana — with the highest densities in the southeastern states. They are animals of edges and margins: agricultural fields, pine flatwoods, abandoned farm buildings, forest clearings, brushy roadsides, and anywhere that supports concentrations of small rodents. In the wild, they are **crepuscular and nocturnal** — most active at dawn, dusk, and after dark, spending the day coiled in rodent burrows, under bark, within wall cavities of old barns, and beneath surface debris. They are **powerful constrictors** — they kill prey by coiling tightly and tightening each time the prey exhales until the prey's circulatory system stops, a method they apply to frozen-thawed prey in captivity with the same efficiency.

The name "corn snake" likely draws from two overlapping sources: their frequent presence in corn-storage barns and granaries, hunting the rodents attracted to stored grain, and the bold checkered black-and-white belly pattern that resembles Indian corn. Their orange and red dorsal saddle markings are occasionally mistaken for the Eastern copperhead by people unfamiliar with the pattern differences — corn snakes have rounded pupils, a slender neck-to-body transition, no facial pit organs, and a distinctly different saddle shape — but corn snakes are entirely non-venomous and pose no biological risk to humans.

Trait	Detail
Common name	Corn snake (also: red rat snake)
Scientific name	<i>Pantherophis guttatus</i>
Native range	Eastern and central USA — New Jersey to Florida, west through Oklahoma and Louisiana
Wild habitat	Agricultural fields, abandoned farm buildings, forest edges, pine flatwoods, brushy margins
Adult length	Typically 3.5 to 5 feet ; occasional individuals reach 5.5 to 6 feet
Adult weight	Typically 250–900 g depending on sex and body condition
Typical lifespan	15–20 years with good care; some captive animals exceed 20 years
Activity pattern	Crepuscular and nocturnal — most active at dawn, dusk, and after dark
Social structure	Strictly solitary — never house together
Temperament	Active, curious, tolerant of handling, rarely defensive once settled; fast and exploratory
Feeding method	Constriction — coils around prey and tightens until prey's heart stops

Color morphs and what they are

Wild corn snakes show the classic pattern: orange-red dorsal saddles with dark borders on a pale orange-gold background, checkered black-and-white belly. Decades of selective breeding in captivity have produced hundreds of **color morphs** — genetic variants that alter pigmentation, pattern, scale structure, or combinations thereof. Common morphs include **amelanistic** (amel — red and white, no black pigment), **anerythristic** (anery — black, gray, and white, no red), **snow** (amel and anery combined — white with pale pink highlights), **okeetee** (wild type with intensified colors and thickened saddle borders), **tessera** (lateral stripe pattern modification), and **scaleless** (reduced scale coverage). Every morph is the same species with identical husbandry requirements. Morph selection is entirely aesthetic preference, not a care variable. Note that hatchlings of some morphs look substantially different from their adult coloration — they can shift significantly in the first twelve to eighteen months as adult pigmentation develops.

What surprises new owners

The activity level is the first genuine surprise. A corn snake does not spend twenty-three hours coiled in a hide the way a ball python often does. She actively explores her enclosure during the evening and night, investigates every branch and corner methodically, and is visibly curious about movement outside the enclosure. This is exactly what you want — a healthy animal doing what corn snakes do — but new owners expecting a sedentary display animal sometimes find the constant movement unexpected and wonder if something is wrong. Nothing is wrong.

The escape artistry is the second surprise, and this one has real consequences. Corn snakes detect temperature differentials through gaps barely larger than their head, and their slender build allows them to fit through spaces that look impossible to the human eye. An enclosure that held a ball python securely may be inadequate for a corn snake who has decided to test every seam systematically. Check every lid every time. Confirm every latch every night. A corn snake found loose in a house after three days has likely been inside a wall cavity or under a large appliance the entire time, and in a cold room that carries real hypothermia risk.

The third surprise is how reliably they eat. New owners who have absorbed ball python forum culture arrive with background anxiety about prolonged food refusals. Corn snakes occasionally decline a meal — briefly, for identifiable reasons — but the months-long unexplained fasting episodes common in ball pythons are not typical of the species. An otherwise healthy corn snake who refuses two or three consecutive meals almost always has a traceable husbandry cause. This is good news, and it means that when a corn snake does refuse food persistently, it deserves prompt investigation rather than the extended "wait and see" patience that is often appropriate for ball pythons.

SECTION 3

Enclosure Setup & Layout

Footprint and the corn snake's use of space

The minimum enclosure for one adult corn snake is 4' × 2' × 2' — the footprint of a 40-gallon breeder or a purpose-built reptile enclosure in the same dimension class. Corn snakes are meaningfully active and use floor length more than height, and the 4-foot floor measurement is the genuine minimum to support normal movement and a complete warm-to-cool gradient. A 5' × 2' × 2' or 4' × 2' × 4' is a legitimate long-term upgrade, and corn snakes will use additional floor length and vertical climbing space. A 5 or 6-foot enclosure for an adult is not excessive — it becomes the snake's entire world for twenty years.

For hatchlings and small juveniles, the concern is the reverse: an enclosure that is too large with too much open, unbroken floor space can suppress feeding. This is not a reason to use a permanently undersized enclosure — it is a reason to furnish a correct-size enclosure thoroughly enough that the snake always has visual cover within a few body lengths. Dense foliage, multiple hides, and cork bark pieces placed across the floor turn a large enclosure into a complex one, which is what a young snake needs.

The thermal gradient

The gradient runs lengthwise from warm end to cool end. A corn snake needs access to warmer temperatures for digestion and immune function, and access to genuinely cooler temperatures for thermoregulation — she should be able to choose any point along the gradient, not oscillate between only two extremes.

Warm side surface temperature: 85–90°F at substrate level. This is warm enough for efficient digestion and measurably cooler than the 88–92°F appropriate for ball pythons, reflecting the corn snake's more temperate native climate. **Warm side ambient air:** 80–85°F at snake height. **Cool end:** 72–78°F — meaningfully cooler than the 76–80°F required for ball pythons and appropriate for the southeastern United States climate that corn snakes evolved in. **Ambient room temperature:** should never drop below 65°F overnight. A room that drops significantly in winter needs a secondary thermostatted heat source or an overnight setback mode on the primary thermostat.

The hides: function and placement

Two hides are the minimum — one warm-side, one cool-side — and both must be **snug-fitting and opaque**. A hide the snake fills almost completely when inside is a hide she will actually use. A hide with excess interior space offers no psychological security. Unlike ball pythons, corn snakes do not spend the entire day in a single hide; they move between hides, perch on branches, and investigate the enclosure throughout the evening and night. A third hide or multiple pieces of cork bark placed at various positions across the enclosure floor increase behavioral complexity and give the snake genuine options throughout the day.

The warm-side hide should be positioned directly over or immediately adjacent to the warmest section of substrate — this is where the snake digests meals and spends the majority of her resting hours. Check the temperature inside the warm hide itself: if it reads significantly below the substrate surface temperature, reposition the hide closer to the heat source.

Visual clutter, climbing, and lid security

Corn snakes move through cover in the wild almost constantly. An unobstructed enclosure is a stressed snake. Arrange the setup so that there is always visual cover within a few body lengths — flat cork, foliage panels, horizontal perches, and positioned hides should break sightlines from end to end. At least one robust climbing branch or piece of driftwood at a shallow-to-moderate incline gives the snake vertical options she will reliably use during her active period. The branch must be firmly anchored so that it cannot shift under the snake's full body weight moving at speed — corn snakes are considerably faster than ball pythons, and an unstable branch will fail.

Lid security cannot be overstated. Every enclosure access point needs a functioning latch or clip. Test the full lid perimeter by pressing gently on different sections. A mesh lid needs a clip on each side, not a single center clip. Sliding glass doors need all locks engaged. Enclosure seams that look sealed but flex under pressure are gaps a corn snake will find. Check the seal after every access, every feeding, and every cleaning.

SECTION 4

Substrate Guide

Substrate selection for corn snakes is primarily about providing a surface that maintains moderate humidity without becoming soggy, supports natural burrowing behavior, and poses no ingestion risk when the snake encounters loose material during feeding.

Substrate	Best for	Humidity performance	Notes
Aspen shavings	Adults and juveniles; the traditional corn snake substrate	Good — maintains 40–60% in most setups; dries at the surface quickly while retaining some moisture at depth	The long-standing standard for corn snakes; spot-cleans cleanly; purchase kiln-dried grade to minimize mold risk; avoid fine dust grades
Coconut fiber (coco coir)	Adults; naturalistic setups	Excellent — holds moisture well; may run above 60% in tightly sealed enclosures and require ventilation adjustment	Slightly messier during feeding than aspen; very forgiving of irregular misting
Cypress mulch	Adults; experienced setups	Very good — retains humidity, drains at the surface; chunky grade preferred	Attractive and naturalistic; fine grades compact over time; always air out before first use to reduce mold risk
Bioactive soil mix (coco fiber + organic topsoil + sphagnum, ~60:30:10)	Adults in permanent naturalistic setups	Excellent, especially with live plants incorporated	Requires a clean-up crew (isopods/springtails); higher initial setup investment; self-maintaining and very low-odour long-term
Sphagnum moss (supplementary zone)	Humid hide insert; shed-cycle supplementation	Outstanding moisture retention	Do not use as full floor substrate; as a damp zone inside a hide it provides a moisture gradient the snake can choose
Paper towel or unprinted newspaper	Hatchlings; quarantine setups; post-treatment observation	Poor — cannot hold humidity; requires active management	Essential for monitoring individual feedings and fecal output during the first weeks with a new snake; zero impaction risk
Cedar shavings	Never, for any reptile	—	Aromatic phenols in cedar are toxic via respiratory exposure; no safe use case at any life stage or concentration
Aromatic pine shavings	Never	—	Same phenol concern as cedar; kiln-dried pine heat-treated specifically for reptile use is a distinct product — confirm the specification before use
Calcium sand or fine play sand	Avoid outright	Very poor	Marketed for reptiles; inappropriate for corn snakes; incidental ingestion risk during feeding; holds essentially no moisture
Gravel or pebble substrate	Never	None	No humidity benefit, significant impaction risk if ingested, injury risk to ventral scales; no valid use case in any snake enclosure

The practical recommendation for most corn snake keepers: kiln-dried aspen shavings at 2–3 inches depth is the cleanest starting point and has been the standard corn snake substrate for decades. It spot-cleans easily, is widely available, holds 40–60% humidity in a standard enclosure without constant active management, and is comfortable for the snake to burrow through. Coconut fiber is an excellent alternative for keepers who prefer a more naturalistic appearance. Bioactive is the long-term upgrade worth moving toward once you understand how your specific enclosure manages its humidity baseline.

SECTION 5

Heat, UVB & Humidity — The 3 Numbers

These three parameters govern the large majority of a corn snake's long-term health. A snake in a correctly heated, correctly humidified enclosure with a stable gradient is a snake whose digestion, immune system, shed cycle, and behavior are all working as they should. A snake in a flat ambient-temperature box at 35% humidity is fighting sub-optimal conditions silently, and the signal will arrive as a patchy shed or a respiratory infection before anything more obvious appears.

Parameter	Target value	How to measure it	What goes wrong when it's off
Warm side surface temperature	85–90°F	Infrared temperature gun at substrate surface; probe thermometer resting on warm-side floor	Too low: poor digestion, sluggish metabolism, persistent feeding refusals; too high: warm-side avoidance, thermal burn risk
Warm side ambient air (snake height)	80–85°F	Probe thermometer 3–4 inches above warm-side floor	Below 78°F: digestion compromised; above 90°F: heat stress, warm-side avoidance
Cool end air temperature	72–78°F	Probe thermometer at cool-end floor level	Below 70°F: metabolic slow-down, immune suppression; above 80°F: no genuine cool zone for thermoregulation
Ambient room temperature	Minimum 65°F overnight	Room thermometer; critical to monitor overnight in winter	Below 60°F: hypothermia risk regardless of enclosure heater performance
UVB (Ferguson zone 1)	UVI 0.5–1.0	Solarmeter 6.5R UV index meter; or manufacturer's published distance chart for the specific bulb model	Absent: likely no measurable harm in most animals, but modest improvement in bone density and natural behavior documented with UVB present; above UVI 2: overexposure risk
Ambient enclosure humidity	40–60% (routine); 60–70% during active shed	Digital hygrometer at mid-enclosure, away from the water dish	Below 35%: dysecdysis (incomplete shedding), retained eye caps, chronic dehydration; above 75% with poor ventilation: scale rot and respiratory infection risk

The UVB question: an honest position


Corn snakes are primarily crepuscular and nocturnal, and many captive animals have lived full, healthy lives without any UVB exposure. This is factually true and worth stating clearly. The current research picture has, however, moved: field studies confirm that corn snakes in the wild encounter ambient UV at dawn and dusk, and that **Ferguson zone 1** exposure — UVI 0.5 to 1.0, roughly equivalent to dappled forest edge light — is associated with modest improvements in D3 synthesis, calcium metabolism, and natural activity patterns in colubrids. A T5 HO 2.0% or low-output 5.0% fluorescent tube at the manufacturer-specified distance provides this level safely on a 12-hour photoperiod. Replace it every 12 months even when it still visibly lights — UV output degrades well before visible light output does. If you choose to keep your corn snake without UVB, dietary D3 supplementation from appropriately dusted prey is a reasonable precaution for a non-breeding adult. The honest frame: UVB probably helps, clearly does not harm a correctly set up enclosure, and the case for including it has grown steadily stronger over the last several years.

Thermostats matched to heat sources

Heat source	Correct thermostat type	Why
Halogen flood lamp	Dimmer thermostat (proportional)	Smoothly reduces voltage to maintain surface temperature without visible flickering
Radiant heat panel (RHP)	Dimmer thermostat	Performs best with smooth voltage regulation; on/off cycling stresses the heating element over time
Ceramic heat emitter (CHE)	Pulse proportional thermostat	No visible light output; pulse-proportional handles pulsed current correctly; dimmer thermostats can damage CHE elements
Under-tank heater (secondary)	On/off thermostat with probe between mat and enclosure floor	Low wattage secondary source; on/off cycling is acceptable here; probe placement between mat and floor prevents surface temperatures exceeding 90°F

SECTION 6

Feeding & Nutrition



Prey sized to the snake's mid-body width, from pinkies for hatchlings to adult mice for a full-grown corn.

Corn snakes are **strict carnivores** whose complete nutritional requirement comes from whole warm-blooded prey. In the wild they are versatile hunters — mice, rats, small birds, bird eggs, and occasionally small lizards all form part of their diet, with small rodents dominant — and they kill everything by **constriction**: coiling around the prey and tightening each time it exhales until the prey's circulatory system stops. In captivity, that hunting drive is entirely intact. A properly warmed frozen-thawed mouse presented with tongs triggers the same behavioral sequence as a wild hunt. The goal of captive feeding is simply to replicate that nutritional transaction safely, hygienically, and on a schedule that matches the snake's actual metabolic needs.

Frozen-thawed and pre-killed only

This guide strongly recommends **frozen-thawed rodents** as the primary and exclusive feeding method. Freshly **pre-killed** prey — dispatched immediately before offering, never frozen — is an acceptable alternative. **Live prey should not be used** under normal circumstances, and this applies to pinky mice as readily as to adult rats.

A live rodent of any size will bite and scratch. Bite wounds to a snake's face and eyes become infected rapidly — a single unsupervised adult rat has blinded and fatally injured healthy snakes. Beyond the injury risk, there is no nutritional advantage to live prey. Frozen-thawed rodents are safer, more hygienic, significantly easier to store in bulk, and accepted by corn snakes — which are typically excellent feeders — just as readily once established on them.

How to thaw and present frozen-thawed prey: place the frozen rodent in a sealed zip-lock bag and submerge it in warm water at approximately 100–105°F. Not hot — hot water cooks the exterior before the interior thaws and produces a temperature inconsistency the snake may detect and reject. Allow 15–20 minutes for a mouse, longer for a small rat, until the prey is fully thawed with no cold spots at the core. Dry the exterior. Confirm the surface is warm to the touch — approximately body temperature. Offer with tongs rather than fingers. Tongs keep your hand outside the strike zone and prevent your fingers from carrying rodent scent after the session.

If your snake was established on live prey before you acquired her and refuses frozen-thawed, the transition is usually simple with corn snakes: offer the warmed frozen mouse in the same spot where the snake has previously struck. Most corn snakes make the switch in one to three attempts. If she still refuses, try **scenting** — lightly rubbing the outside of the frozen prey with a live prey item briefly brought into contact and then removed before offering. This scenting technique is rarely necessary with corn snakes but is almost always effective when it is.

Prey sizing

Prey should be **no wider at its thickest point than the snake's body at its thickest point** — roughly the mid-body girth. When in doubt, size down rather than up. A prey item slightly smaller than the maximum is always safer than one that is too large. Oversize prey dramatically increases regurgitation risk, and a corn snake who has regurgitated should not be offered food again for at least ten to fourteen days to allow the digestive system to recover fully.

Snake life stage	Approximate length	Appropriate prey type
Hatchling	Under 12 inches	Pinky mouse (newborn, unfurred)
Young juvenile	12–18 inches	Fuzzy mouse (lightly furred)
Growing juvenile	18–24 inches	Hopper mouse (weaned)
Sub-adult	24–36 inches	Small adult mouse
Adult	36–48 inches	Adult mouse; large-bodied adults may progress to small rats
Large adult	48 inches and over	Adult mouse or small rat, matched to current mid-body girth

Feeding schedule

Life stage	Age	Frequency
Hatchling	0–6 months	Every 5–7 days
Juvenile	6–18 months	Every 7–10 days
Sub-adult	18 months–3 years	Every 10–14 days
Adult	3 years and older	Every 10–14 days

Feeding strikes: what to do

Corn snakes have a considerably better feeding record than ball pythons. Prolonged, open-ended fasting of weeks or months without an identifiable cause is uncommon in a healthy, correctly kept corn snake. When a corn snake refuses food, the cause is almost always traceable.

Common causes to check first: the snake is in the pre-shed blue phase (wait for the shed to complete and give it a few days); the thermal gradient has drifted below the correct warm-side temperature (measure it with fresh probe readings); the snake is reacting

to recent handling; humidity is below 35%; or the snake is showing a mild seasonal cycling response in late autumn. Some corn snakes reduce feeding drive briefly in November through January — this is usually one or two refusals, not four months of total abstinence.

What to do during a feeding refusal: verify the gradient with fresh readings; confirm humidity is in range; offer prey at the snake's most active time — dusk or after dark; try a prey item one size smaller; try brief scenting with a different prey species; offer prey inside a small paper bag or cardboard box, which sometimes triggers a more confident strike in a hesitant snake; eliminate all handling during the refusal period and give ten days between attempts. **When to call a vet:** persistent refusal across three or more consecutive offerings with no identifiable cause, any weight loss visible as a thinning body cross-section, or any concurrent physical change — mucus, wheezing, mouth changes, unusual lethargy during active hours.

***Shop the food and supplies:** Browse Reptile & Amphibian Food and Live Feed Delivered!.*

SECTION 7

Shedding

A healthy corn snake sheds her entire skin in a single unbroken piece every **four to eight weeks** as a growing juvenile and every **six to twelve weeks** as an adult. Frequency varies with growth rate, feeding regularity, season, and individual variation. Shedding is normal, is not painful, and requires no intervention from you if enclosure humidity is correctly maintained. Your job is to observe the shed and treat any problem the shed reveals as information about the enclosure conditions that need correcting.

The shed cycle

The first visible sign of an impending shed is the **blue phase**: the eyes shift from clear and sharp to a milky, opalescent gray-blue as a fluid layer accumulates between the old and new skin at the eye surface. Corn snakes in the blue phase have significantly impaired vision, typically refuse food, may become more defensive, and often spend additional time near any humid zone in the enclosure. Do not offer food during the blue phase. Handle only if strictly necessary, and approach slowly from the side. After three to five days, the eyes clear — the fluid has redistributed, but the shed itself is still two to four days away. Increase enclosure humidity and check it daily once the eyes clear.

The shed itself typically happens at night. The snake rubs her nose tip on a rough surface — a branch, a hide entrance, a piece of cork bark — to initiate a lip tear, then works the old skin backwards off her body as a continuous inside-out tube. A complete, single-piece shed that includes intact **eye caps** — the transparent, permanently fused scale called the **spectacle** or **brille** that covers the eye in place of eyelids — confirms that enclosure humidity and hydration are both correct.

Good and bad sheds

A good shed comes off in a single continuous piece or two large sections. The eye caps are visible as small clear domes in the head portion of the shed skin. The shed feels intact and slightly stretchy rather than dry and papery. A bad shed fragments into many small pieces, leaves gray patches of retained skin on the body once it dries, or is missing one or both eye caps — meaning they may still be on the snake's eyes.

Retained eye caps are the most clinically serious shed complication. A spectacle that did not release with the rest of the head skin appears as a slightly thickened, dull, subtly textured layer over one or both eyes. Left in place through two or more additional shed cycles, retained spectacles build up and eventually press on the eye surface, causing permanent damage including partial or complete blindness.

What to do for stuck shed: increase enclosure humidity to 65–70% for 24–48 hours. Offer a shallow warm soak — belly-depth water, head free, at approximately 85°F — for 20 to 30 minutes. After soaking, run damp fingers gently along the body; softened retained shed typically releases with minimal pressure. Never peel dry retained shed — it is adhered to the fresh skin beneath, and pulling dry shed risks tearing the new scale layer and leaving raw tissue exposed.

What to do for a retained eye cap: the soak-first rule is absolute. After a full warm soak, place a lightly damp paper towel in the enclosure and let the snake move across it — the slight surface texture combined with a softened spectacle sometimes produces a release that the soak began. Do not attempt to pull a retained eye cap with dry fingers or any tool; the margin for eye damage is small. If two complete soaking sessions do not free the cap, this is a veterinary appointment.

SECTION 8

Common Challenges & Solutions

Feeding strike

What you see: the snake investigates prey with her tongue and retreats; does not strike; ignores the prey entirely; or refuses to emerge from the hide when prey is offered. **Why:** most commonly pre-shed, a thermal gradient that has drifted out of range, low humidity, recent handling, a recent enclosure move, or a mild seasonal cycling response in late autumn or winter. Less commonly, early illness. **What to do:** run through the husbandry checklist — measure temperatures with fresh probe readings, check humidity, review handling in the last 48 hours, check for signs of pre-shed. Eliminate all handling during the refusal period. Wait ten days and try again. **When to call a vet:** three or more consecutive refusals with no identifiable husbandry cause, any concurrent physical symptom, or visible muscle wasting.

Respiratory infection

What you see: audible wheezing, clicking, or crackling during normal breathing; mucus or bubbles at the nostrils or inside the mouth; the snake holding her head slightly elevated; visibly labored breathing with exaggerated body movement. **Why:** bacterial or viral airway infection, most often triggered by chronically low ambient temperatures, sustained high humidity combined with poor ventilation, or immune suppression from prolonged enclosure stress. **What to do:** correct any temperature deficiency immediately — a respiratory infection advances much faster in a cold snake than a warm one. Isolate from any other reptiles in the household. **When to call a vet:** within one to two days of noticing any respiratory symptom. These infections respond well to appropriate antibiotics when caught early and become serious quickly when they are not.

Snake mites

What you see: small, dark, moving specks — approximately the size of a printed period — on the snake's skin surface, floating in the water dish, and accumulating in enclosure corners. The snake may soak for unusual lengths of time or rub against surfaces. Mites concentrate under chin scales, around the eye area, and in scale folds along the spine. **Why:** **snake mites** (*Ophionyssus natricis*) are the most common external parasite of captive snakes, reproduce rapidly, and are typically introduced through a new animal from an expo or store, second-hand equipment, or close proximity to an infested collection. **What to do:** quarantine immediately. Remove and discard all substrate and porous decor. Scrub all enclosure surfaces with a reptile-safe disinfectant. Treat with a reptile-approved mite product following label timing exactly, including a mandatory second application on schedule to catch mites that hatch after the first treatment. Soak the snake in lukewarm water for 15–20 minutes during enclosure treatment to drown surface mites. Wash your hands and clothing thoroughly before and after handling the snake. **When to call a vet:** heavy infestation, a hatchling or recently unwell snake, or no response after two correctly timed applications.

Dysecdysis (stuck or incomplete shed)

What you see: gray, papery retained skin patches on the body after a shed; the shed arriving in fragments rather than a single piece; retained skin constricting the tail tip; retained eye cap(s) visible as a thickened or dull layer over one or both eyes. **Why:** enclosure humidity consistently below 40% during the shed cycle, chronic dehydration from a dirty or unavailable water dish, or no rough surface provided for the snake to initiate the lip tear. **What to do:** increase humidity to 65–70% for 24–48 hours; warm shallow soak for 20–30 minutes; gentle assistance with damp fingers along the body. For a retained eye cap, follow the soak with a damp paper towel pass. Never peel dry retained skin — always soak first. **When to call a vet:** retained eye cap not releasing after two soaking sessions; skin constricting the tail tip with any sign of circulation compromise visible as darkening or swollen tissue distal to the constriction.

Thermal burns

What you see: blistered, lifted, or discolored scales on the belly or sides; the snake actively avoiding the warm end of the enclosure; in more serious cases, exposed tissue beneath lifted scales. **Why:** contact with an unregulated heat source — almost always a heat rock, an unthermostated under-tank mat, or any heat source the thermostat probe has been positioned too far from to read accurately. Corn snakes, like all reptiles, have a slower pain response than mammals and may not self-regulate off a dangerously hot surface before damage occurs. **What to do:** remove or immediately correct the unregulated source; transfer the snake to a clean, thermostatted enclosure on paper towel substrate; keep the wound dry and free from substrate particles. **When to call a vet:** any burn with exposed or necrotic tissue, any burn near the cloaca, or any wound showing signs of spreading infection.

Mouth rot (infectious stomatitis)

What you see: the snake rubbing her mouth against enclosure walls or decor; swelling or redness in the gum tissue; in more advanced cases, yellow or cheese-like discharge at the mouth margins; reduced interest in food. **Why:** bacterial infection of the oral mucosa, typically initiated by a minor bite from frozen-thawed prey that was too large, an abrasion from a rough decor surface, or immune suppression from sustained incorrect temperatures. **What to do:** examine the mouth under good lighting; correct any environmental causes immediately; in mild early cases a dilute chlorhexidine solution applied with a cotton swab can provide some initial relief. **When to call a vet:** within a few days of noticing any mouth tissue change — **infectious stomatitis** spreads to underlying bone quickly and requires antibiotic treatment to resolve.

Constipation and impaction

What you see: no stool for three to four weeks after feeding; a visibly swollen or lumpy lower abdomen; lethargy; in severe cases, visible straining at the cloaca with no result. **Why:** a prey item that was too large, chronic dehydration, incidental substrate ingestion during feeding on loose aspen or cypress, or less commonly a foreign object. **What to do:** warm shallow soaks once or twice daily for 15–20 minutes; ensure the water dish is clean and accessible; gentle belly massage from mid-body toward the cloaca after each soak. Use a paper towel or tile as a feeding surface if you suspect the snake is ingesting substrate during feeding. **When to call a vet:** no result after 48 hours of consistent soaking; noticeably hardened or distended abdomen; visible distress at the cloaca.

Escape and systematic recovery

What you see: the corn snake is not in the enclosure. **Why:** one unlocked clip, one latch not fully engaged, one gap in the lid seal that the snake located, tested, and eventually pushed through — corn snakes investigate every enclosure edge methodically on a nightly basis. **What to do:** close and lock the room immediately to contain the search area. Corn snakes seek warmth and tight spaces — search from the floor level upward, checking near heating appliances, behind the refrigerator and large furniture, under baseboards and door thresholds, in closet corners, clothing piles, and inside any open boxes or furniture cavities. Leave the now-empty enclosure open with a warm hide inside and a warmed-prey scent trail nearby — escapees frequently return to a familiar heat source once hungry. Damp paper towels placed flat on the floor overnight will show crossing marks from the snake's ventral scales. In winter or in cool rooms, a snake who has been loose for more than a day will be cold and sluggish. **When to call a vet:** if found significantly dehydrated, exhibiting cold-stunning symptoms, or injured.

Regurgitation

What you see: the snake expels a partially or fully intact meal — typically as a mucus-coated cylinder of roughly prey size — hours to days after feeding. **Why:** most commonly caused by handling within 48 hours of feeding, prey that was too large, prey that was not fully thawed before offering, the thermal gradient dropping too low after feeding prevented proper digestion, or a significant temperature stress event. Less commonly, an early sign of gastrointestinal infection. **What to do:** do not offer food again for at least 10–14 days — the digestive system needs recovery time after a regurgitation. On the next offering, use prey one size smaller, confirm the warm side is maintaining the correct temperature, and ensure the snake is not handled for at least 48 hours before or after. **When to call a vet:** two consecutive regurgitation events, or regurgitation accompanied by any other physical symptom.

Obesity from overfeeding

What you see: an almost circular cross-section when viewed head-on; visible fat deposits along the flanks and immediately behind the head; sluggish, reluctant movement during the normally active evening period. **Why:** corn snakes are metabolically efficient animals adapted to periodic hunting — they are not designed for consistently available prey on a short interval. Overfeeding is common, especially when a snake accepts food enthusiastically at every offering. Chronic obesity in corn snakes is associated with fatty liver disease, reproductive complications in females, and reduced lifespan. **What to do:** reduce prey size before reducing feeding frequency; size prey accurately to the snake's actual current mid-body girth rather than the maximum she will accept. **When to call a vet:** a severely obese snake, or one refusing all food for more than six weeks during a diet correction.

Quick reference

What you see	Likely cause	What to do	Vet urgency
Food refused; active at dusk, maintaining weight	Pre-shed, husbandry drift, mild seasonal slow-down	Check gradient and humidity; wait 10 days; no handling	Only if 3+ consecutive refusals or muscle wasting
Wheezing, mucus at nostrils, head elevated	Respiratory infection	Correct temperatures; isolate	Within 1–2 days
Moving specks on skin and in water dish, soaking excessively	Snake mites	Quarantine; treat enclosure and snake	Heavy infestation or hatchling: soon
Patchy retained gray skin; dull or thickened eye surface	Dysecdysis / retained eye cap	Warm soak 20–30 min; damp paper towel surface	Retained eye cap after 2 soaks: same week
Blistered or lifting belly scales	Thermal burn	Remove unregulated source; paper towel substrate	Exposed or necrotic tissue: same day
Mouth wiping, redness or swelling at mouth margins	Mouth rot / infectious stomatitis	Chlorhexidine rinse; correct stressors	Within a few days
No stool 3–4+ weeks after feeding, swollen abdomen	Constipation / impaction	Warm soaks; gentle belly massage	48 hours with no result
Snake not in enclosure	Escape	Systematic room search; leave warm enclosure open with hide	If found dehydrated, cold, or injured
Prey expelled hours to days post-feeding	Regurgitation	10–14 day fast; next prey item smaller; no handling 48 hrs pre/post	Two consecutive events
Circular cross-section; fat deposits along flanks	Obesity from overfeeding	Reduce prey size; enforce correct schedule	Refusing all food 6+ weeks during correction

SECTION 9

Handling

Corn snakes are among the most naturally handleable snakes in captivity. A corn snake raised with consistent, calm handling from a young age is typically alert, responsive, and genuinely non-defensive as an adult. The key word is consistent: a snake who is handled occasionally and unpredictably is more reactive than one whose routine includes regular positive contact. Handling frequency of two to three times per week for 10 to 15 minutes per session produces a settled, confident adult. More than that is fine; less than once a week keeps the snake in a slightly higher state of alertness during sessions.

How to pick up a corn snake

Approach from the side or from below — never from above. A hand descending from above mimics a predator; it is exactly what the corn snake's aerial threat response is calibrated for, and it will reliably produce an S-curve posture in a snake who would otherwise be perfectly calm approached from the side. Slide your primary hand under the front third of the snake's body and bring your second hand under the rear portion to support the full length. Lift slowly and evenly. Corn snakes move faster than ball pythons and begin exploring immediately — let them "treadmill" forward from hand to hand rather than gripping them still. A snake who is moving is a snake who is not coiling into a defensive posture. Support the full body length at all times; a snake whose rear third hangs unsupported will coil around your forearm reflexively.

Reading warning signs

Corn snakes communicate clearly when they want the interaction to end.

S-curve posture: the snake draws her neck and anterior body into a lateral S with the head pulled back and elevated. This is the pre-strike posture — she is loaded. It is common in young hatchlings who are still establishing trust with their keeper and diminishes with consistent positive handling. When you see it, set her down calmly and end the session without abrupt movement.

Tail rattling: corn snakes can vibrate their tail tip against substrate — a behavioral mimicry of rattlesnakes that is entirely harmless but worth recognizing as a warning signal. End the session when this occurs.

Musking: the snake releases a pungent secretion from cloacal glands. Harmless, unpleasant, and common in juveniles during their first weeks of regular handling. It diminishes substantially as the snake settles with consistent contact. Wash your hands after the session.

A bite: a defensive bite from a young corn snake is a row of small puncture wounds. It is not dangerous, mildly uncomfortable, and not a reason to stop working with the snake long-term. It is a reason to examine what triggered the response and adjust your approach for the next session.

When NOT to handle

- **Within 48 hours of feeding** — handling too soon after feeding is the most common cause of regurgitation, which is stressful for the snake and carries digestive health risks on re-ingestion.
- **During the blue phase** of the shed cycle — vision is severely impaired and defensive responses are elevated; the additional stress is not useful.
- **During a feeding strike or food refusal** — adding handling stress to an already stressed animal compounds the problem rather than resolving it.
- **For the first five to seven days after arrival** in a new enclosure — allow the snake to map the space and establish it as safe before handling sessions begin.

Children

Corn snakes are well-suited to gentle children under close adult supervision. Have children handle seated on the floor — never standing — with an adult's hands directly underneath as an immediate safety net. Teach children to let the snake move forward rather than grip or constrict her movement. A corn snake who begins to S-curve should be calmly returned to her enclosure without alarm.

SECTION 10

Routine Care Calendar

How often	What to do
Daily	Check the water dish — full and clean; replace immediately if fouled
Daily	Verify warm-side and cool-end temperatures with probe thermometers
Daily	Check enclosure humidity; mist if below 40% in routine conditions or below 60% during an active shed cycle
Daily	Spot-clean — remove visible feces, urates, regurgitated material, or shed fragments
Daily	Observe the snake at dusk: active? Alert? Normal posture and movement?
Daily	Confirm the lid is fully secured with all clips or locks engaged
Every 5–14 days	Offer food on the correct schedule for life stage; log whether the offering was accepted or refused
Weekly	Replace and scrub the water dish in hot water with mild unscented soap; dry fully before returning
Weekly	Weigh the snake on a kitchen scale; record and compare to the previous week
Weekly	Inspect the snake's full body during or after a handling session for retained shed, scale changes, wounds, or swelling
Weekly	Check all enclosure seals and lid hardware; confirm every latch is functional
Monthly	Deep clean — remove all decor; scrub all interior surfaces with a reptile-safe disinfectant; dry completely before replacing substrate and furnishings
Monthly	Partial or full substrate replacement depending on the system; spot-clean soiled areas promptly rather than waiting for monthly changes
Monthly	Photograph the snake from above for a body-condition baseline to compare over time
Monthly	Review the food and weight log; note any trend in mass or feeding acceptance rate
Every 6 months	Audit the thermal gradient with fresh probe thermometer batteries; replace any probes giving inconsistent or drifted readings
Every 12 months	Replace the UVB bulb regardless of visible light output — UV output degrades before visible light output does
Annually	Exotic reptile vet wellness check including a fecal exam for internal parasites; twice annually for breeding females or snakes over 12 years old

SECTION 11

When to See a Vet

Corn snakes are hardy animals who rarely show illness until a condition is already well established. The rule: **when you are not sure whether to call, call.**

Go today — same-day or emergency:

- Any prolapse of tissue from the cloaca — visible tissue that cannot be retracted
- Open-mouth breathing, or any audible wheezing, clicking, or crackling during respiration
- A snake who is unresponsive, completely limp to gentle touch, or cannot right herself
- Wound with exposed tissue or bleeding that does not slow within a few minutes
- Thermal burn with exposed or visibly necrotic tissue
- A retained eye cap that appears to be pressing on or distorting the eye surface
- Any suspected outdoor exposure to pesticides, toxic plants, or venomous species if the snake escaped outside

Book within a few days:

- No stool for three or more weeks with no response to warm soaking
- Retained shed or eye cap not releasing after two complete soaking sessions
- Progressive muscle wasting — body cross-section shifting from rounded to angular
- Mouth tissue that is red, swollen, or showing any discharge
- Respiratory sounds that appeared recently and have not resolved in 24–48 hours
- Any new firm lump anywhere on the body
- Two consecutive regurgitation events with no obvious environmental cause
- Persistent food refusal across three or more consecutive offerings with no identifiable husbandry reason

Find your exotic reptile vet now, before you need one. Confirm they see colubrids specifically — corn snakes are common enough that most exotic practices see them, but confirm rather than assume. Bring a temperature log, a feeding record, a weight history, and a fresh fecal sample in a sealed container.

SECTION 12

FAQ

What do corn snakes eat?

Corn snakes are strict carnivores and eat exclusively whole warm-blooded prey. In the wild they hunt small rodents — primarily mice and rats — along with small birds, bird eggs, and occasionally small lizards, killing all prey by constriction. In captivity, frozen-thawed or freshly pre-killed mice and small rats are the correct and strongly preferred food source. Live prey should never be used under normal circumstances: a live rodent will bite and scratch, bite wounds to a snake's face and eyes become infected rapidly, and there is no nutritional benefit that justifies the risk. Prey is sized to the snake's mid-body girth — hatchlings start on pinky mice and progress to adult mice and occasional small rats as adults. Corn snakes do not eat insects, fish, fruit, or any plant material.

How big do corn snakes get?

Adult corn snakes typically reach **3.5 to 5 feet** in length and weigh between 250 and 900 grams depending on sex, genetics, and body condition. Females tend to be slightly larger than males on average, though the difference is less pronounced than in ball pythons. Occasional individuals reach 5.5 to 6 feet, but animals of that size are outside the typical range. Most of the visible growth occurs in the first two to three years; after that, growth continues slowly and imperceptibly. The slender, lightweight build makes even a 5-foot corn snake straightforward to handle and support compared to a stockier species of similar length.

How long do corn snakes live?

With good care, **15 to 20 years** is the realistic expectation for a captive corn snake, and well-kept individuals sometimes exceed 20 years. The factors most consistently associated with reaching the upper end of that range are correct temperatures maintained throughout the snake's life, humidity kept in the 40–60% range, a feeding protocol that avoids chronic overfeeding, and regular veterinary care that catches problems while they are still minor. This is a fifteen to twenty year commitment — longer than most people own a car — and that deserves to be taken seriously before the first purchase.

What size enclosure does a corn snake need?

The minimum for one adult corn snake is **4 feet × 2 feet × 2 feet**. Corn snakes are active animals who use floor length meaningfully, and the 4-foot minimum is a genuine floor, not a conservative suggestion. A 5' × 2' × 2' or larger enclosure is a legitimate long-term upgrade that a corn snake will make full use of. Front-opening or side-opening enclosures are preferred over top-opening tanks because an overhead hand mimics a predator. Whatever dimensions you choose, the enclosure must seal completely and latch securely — corn snakes are expert escape artists, and a gap that seems negligible is a gap they will locate and exploit.

How often do I feed my corn snake?

Hatchlings eat every 5 to 7 days; juveniles every 7 to 10 days; adults every 10 to 14 days. Corn snakes are metabolically efficient animals not built for frequent, consistently available meals — overfeeding on a short interval leads to the obesity that is increasingly common in captive corn snakes. Feed on the correct schedule with correctly sized prey, and resist offering extra meals when the snake accepts eagerly. A corn snake who consistently accepts food on the correct schedule is doing exactly what a healthy corn snake should do, and that reliable acceptance is not an invitation to feed more often.

Do corn snakes need UVB light?

UVB is beneficial but not strictly required for corn snakes. Many captive corn snakes have lived full, healthy lives without any UVB exposure, and this is worth stating honestly. Current research indicates that low-level UVB — Ferguson zone 1 exposure providing UVI 0.5 to 1.0, equivalent to dappled forest edge light — offers modest improvements in D3 synthesis, calcium

metabolism, and natural activity patterns without meaningful risk in a correctly set up enclosure. A low-output T5 HO fluorescent tube on a 12-hour photoperiod achieves this safely. Replace it annually, because UV output degrades significantly before visible light output does. If you choose to keep your corn snake without UVB, appropriately D3-supplemented prey is a reasonable precaution. The honest position: UVB probably helps, clearly does not harm a correctly maintained enclosure, and the evidence for including it has grown consistently stronger over the last decade.

What humidity does a corn snake need?

Corn snakes need **40 to 60% relative humidity** as a routine baseline, raised to **60 to 70% during active shed cycles**. This is meaningfully lower than ball pythons and reflects the drier habitats across the eastern United States where corn snakes originate. Humidity below 35% causes dysecdysis — incomplete shedding — and chronic dehydration. Humidity consistently above 75% combined with poor ventilation creates conditions for scale rot and respiratory infection. Most standard setups achieve the 40–60% baseline naturally with aspen shavings at 2–3 inches depth, a clean water dish, and a reasonably sealed enclosure. A hygrometer tells you where your specific setup actually sits rather than where you assume it does.

Why won't my corn snake eat?

The most common reasons, in order of frequency: the snake is in the pre-shed blue phase — wait for the shed to complete before offering food again; the thermal gradient has drifted below the correct warm-side temperature — measure it with fresh probe readings; the snake is reacting to recent handling or a recent move; or there is a mild seasonal slow-down in late autumn or early winter. Unlike ball pythons, corn snakes do not typically go on extended, open-ended fasting episodes without an identifiable cause. A healthy corn snake who refuses two or three consecutive meals almost always has a traceable husbandry reason. Check the gradient, check the humidity, rule out pre-shed, and eliminate all handling during the refusal period. If she is active at her normal evening time and maintaining her body weight, the cause is almost certainly environmental and correctable.

Are corn snakes good for beginners?

Yes — genuinely so, not just as a marketing claim. Corn snakes are widely considered the best beginner snake in the hobby because they accept handling readily, eat reliably without the prolonged unpredictable fasting episodes common in ball pythons, are appropriately sized for most living spaces, and are robust enough to do well once husbandry is correctly established. Their care requirements are straightforward: a correct thermal gradient, moderate humidity at 40–60%, correctly sized frozen-thawed prey on a sensible schedule, and an escape-proof enclosure. None of that is advanced. A corn snake set up correctly and handled consistently is among the most rewarding snakes available at any experience level, and a well-kept individual can be a manageable, curious companion for fifteen to twenty years.

Can two corn snakes live together?

No. Corn snakes are solitary animals in the wild and gain nothing from the presence of another snake in captivity. Housed together, they compete for hides, basking positions, and food — cohabitation always produces a dominant animal and a subordinate whose chronic stress is easy to miss on a casual daily check. Two snakes who appear to be resting peacefully are almost always in a dominant-subordinate relationship; the subordinate loses weight, develops immune problems, and sheds poorly over weeks and months rather than showing immediate distress. There is also a genuine cannibalism risk when similarly sized snakes are fed in a shared enclosure — corn snakes belong to the rat snake family and are known to take other snakes as prey in the wild. One corn snake per enclosure is the standard, it exists for good reason, and there are no exceptions to it.

Ready to shop? Start with *Reptiles & Amphibians*. 10% of every purchase supports no-kill animal shelters.
Not sure this is the right pet? Compare your options in our *Reptile Guide*.

SECTION 13

Every Purchase Gives Back

Before we earn a dime, **10% of every purchase at Walnut Creek Supply goes to no-kill animal shelters across the country.** Not as a checkbox at checkout. Not as an optional donation round-up. Off the top of every order, every day — because responsible pet ownership and community care are not separate decisions.

Corn snakes are surrendered to rescues in significant numbers every year. Most arrive in ten-gallon tanks on calcium sand with an unthermostated heat rock, malnourished and dysecdysic, the survivors of well-intentioned setups built on outdated information that has been on the internet since the early 2000s and has not been updated. They are remarkably resilient animals — given correct temperatures, appropriate humidity, and frozen-thawed prey on a proper schedule, most recover fully. If you are reading this guide, you are already doing the work that animals in rescues never had done for them. And if you can adopt rather than shop — reptile rescues in every region have healthy, handleable corn snakes of all morphs waiting for a keeper who knows what they are doing.

We are a family business in Christiana, Lancaster County, Pennsylvania. We grow our own food, keep our own animals, and stock only what we would put in front of our own pets — **no mystery ingredients**, no heat rocks, no calcium sand sold as a substrate. If you have a question about your corn snake's thermal gradient or her shed cycle, call us at **(717) 983-8069**. We would rather talk you through it.

Welcome to fifteen to twenty years of a snake who knows your scent and comes to the front of her enclosure when you sit beside her.

From our family to yours, Walnut Creek Supply — Christiana, Pennsylvania walnutcreeksupply.com